

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
6 May 2004 (06.05.2004)

PCT
WO 2004/037789 A2

(51) International Patent Classification: C07D 213/00

Darmstadt (DE), FINSINGER, Dietl (DE/DE); Pippinger-
strasse 33, 81245 München (DE).

(21) International Application Number: PCT/EP2003/011134

(74) Common Representative: MERCK PATENT GMBH,
Frankfurter Strasse 250, 64293 Darmstadt (DE).

(22) International Filing Date: 8 October 2003 (08.10.2003)

(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MY, NZ, OM, PG, PH, PL, PT, RO, RU, SC,
SD, SE, SG, SK, SL, SY, TH, TM, TN, TR, TT, TZ, UA,
UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date:
24 October 2002 (24.10.2002) EP
02023906.7
28 July 2003 (28.07.2003) US
60/490,285

(71) Applicant (for all designated States except US): MERCK
PATENT GMBH (DE/DE); Frankfurter Strasse 250,
64293 Darmstadt (DE).

(72) Inventors; and
Inventors/Applicants (for US only): BUCHSTALLER,
Hans-Peter (AT/DE); Heinrichstrasse 54, 64331 Weiter-
stadt (DE); WIESNER, Matthias (DE/DE); Beethoven-
ring 10, 64342 Seelheim-Jugenheim (DE); SCHATZ,
Oliver (DE/DE); Eschenstrasse 32, 63517 Rödertbach
(DE); AMENDT, Christine (DE/DE); Backhausstrasse
22, 64289 Darmstadt (DE); ZENKER, Frank (DE/DE);
Schulzengeisse 7, 64291 Darmstadt (DE); SIRENBERG,
Christian (DE/DE); Tannusstrasse 10, 64289 Darmstadt
(DE); GRIEL, Matthias (DE/DE); Lindenweg 44, 64291

Published:
— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guide
to the Preparation of the International Publication" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: METHYLENE UREA DERIVATIVES

(57) Abstract: The present invention relates to methylene urea derivatives of formula (I), the use of the compounds of formula (I) as
inhibitors of raf-kinase, the use of the compounds of formula (I) for the manufacture of a pharmaceutical composition and a method
of treatment, comprising administering said pharmaceutical composition to a patient.

Methylene urea derivatives

The present invention relates to methylene urea derivatives, methylene urea derivatives as medicaments, methylene urea derivatives as inhibitors of raf-kinase, the use of methylene urea derivatives for the manufacture of a pharmaceutical, a method for producing a pharmaceutical composition containing said methylene urea derivatives, the pharmaceutical composition obtainable by said method and a method of treatment, comprising administering said pharmaceutical composition.

10 Protein phosphorylation is a fundamental process for the regulation of cellular functions. The coordinated action of both protein kinases and phosphatases controls the levels of phosphorylation and, hence, the activity of specific target proteins. One of the predominant roles of protein phosphorylation is in signal transduction, where extracellular signals are amplified and propagated by a cascade of protein phosphorylation and dephosphorylation events, e.g. in the p21^{ras}/raf pathway.

15 The p21^{ras} gene was discovered as an oncogene of the Harvey (ras^H) and Kirsten (ras^K) rat sarcoma viruses. In humans, characteristic mutations in the cellular ras gene (c-ras) have been associated with many different types of cancers. These mutant alleles, which render Ras constitutively active, have been shown to transform cells, such as the murine cell line NIH 3T3, in culture.

25 The p21^{ras} oncogene is a major contributor to the development and progression of human solid cancers and is mutated in 30 % of all human cancers (Bolton et al. (1994) Ann. Rep. Med. Chem., 29, 165-74; Bos. (1989) Cancer Res., 49, 4682-9). Oncogenic Ras mutations have been identified for example in lung cancer, colorectal cancer, pancreas, thyroid cancer, melanoma, bladder tumors, liver tumor, kidney tumor, dermatological tumors and haematological tumors (Doljei et al. (2001), J.

BEST AVAILABLE COPY

Natl. Cancer Inst. 93(14), 1062-74; Midgley, R.S. and Kerr, D.J. (2002) *Critical Rev. Onc/ hematol* 44, 109-120; Downward, J. (2003), *Nature reviews* 3, 11-22). In its normal, unmutated form, the ras protein is a key element of the signal transduction cascade directed by growth factor receptors in almost all tissues (Avruch et al. (1994) *Trends Biochem. Sci.*, 19, 279-83).

5

Biochemically, ras is a guanine nucleotide binding protein, and cycling between a GTP-bound activated and a GDP-bound resting form is strictly controlled by ras endogenous GTPase activity and other regulatory proteins. The ras gene product binds to guanine triphosphate (GTP) and guanine diphosphate (GDP) and hydrolyzes GTP to GDP. It is the GTP-bound state of Ras that is active. In the ras mutants in cancer cells, the endogenous GTPase activity is alleviated and, therefore, the protein delivers constitutive growth signals to downstream effectors such as the enzyme raf kinase. This leads to the cancerous growth of the cells which carry these mutants (Magnuson et al. (1994) *Semin. Cancer Biol.*, 5, 247-53). The ras proto-oncogene requires a functionally intact c-raf1 proto-oncogene in order to transduce growth and differentiation signals initiated by receptor and non-receptor tyrosine kinases in higher eukaryotes.

15

20

Activated Ras is necessary for the activation of the c-raf-1 proto-oncogene, but the biochemical steps through which Ras activates the Raf-1 protein (Ser/Thr) kinase are now well characterized. It has been shown that inhibiting the effect of active ras by inhibiting the raf kinase signaling pathway by administration of deactivating antibodies to raf kinase or by co-expression of dominant negative raf kinase or dominant negative MEK also called ERK, the substrate of raf kinase, leads to the reversion of transformed cells to the normal growth phenotype see: Daum et al. (1994) *Trends Biochem. Sci.*, 19, 474-80; Fridman et al. (1994) *J Biol. Chem.*, 269, 30105-8. Kolch et al. (1991) *Nature*, 349, 426-28) and for review Weinstein-Oppenheim et al. *Pharm. & Therap.* (2000), 88, 229-279.

30

Similarly, inhibition of raf kinase (by antisense oligodeoxynucleotides) has been correlated in vitro and in vivo with inhibition of the growth of a variety of human tumor types (Monia et al., *Nat. Med.* 1996, 2, 668-75; Geiger et al. (1997), *Clin. Cancer Res.* 3(7): 1179-85; Lau et al. (2002), *Antisense Nucl. Acid. Drug Dev.* 12(1): 11-20; McPhillips et al. (2001), *Br. J. Cancer* 85(11): 1753-8).

5

Raf serine- and threonine-specific protein kinases are cytosolic enzymes that stimulate cell growth in a variety of cell systems (Rapp, U.R., et al. (1988) in *The oncogene handbook*; T. Curran, E.P. Reddy, and A. Skalka (ed.) Elsevier Science Publishers; The Netherlands, pp. 213-253; Rapp, U.R., et al. (1988) *Cold Spring Harbor Sym. Quant. Biol.* 53:173-184; Rapp, U.R., et al. (1990) *Inv Curr. Top. Microbiol. Immunol.* Potter and Melchers (eds), Berlin, Springer-Verlag 166:129-139).

10

15

Three isozymes have been characterized:

c-Raf (also named Raf-1, c-raf-1 or c-raf1) (Bonner, T.I., et al. (1986) *Nucleic Acids Res.* 14:1009-1015), A-Raf (Beck, T.W., et al. (1987) *Nucleic Acids Res.* 15:595-609), and B-Raf (Okawa, S., et al. (1998) *Mol. Cell. Biol.* 8:2651-2654; Sithanandam, G. et al. (1990) *Oncogene*:1775).

20

These enzymes differ in their expression in various tissues. Raf-1 is expressed in all organs and in all cell lines that have been examined, and A- and B-Raf are expressed in urogenital and brain tissues, respectively (Storm, S.M. (1990) *Oncogene* 5:345-351).

25

Raf genes are proto-oncogenes: they can initiate malignant transformation of cells when expressed in specifically altered forms. Genetic changes that lead to oncogenic activation generate a constitutively active protein kinase by removal or interference with an N-terminal negative regulatory domain of the protein (Heidecker, G., et al. (1990) *Mol. Cell. Biol.* 10:2503-2512; Rapp, U.R., et al. (1987) in *Oncogenes and cancer* S. A. Aaronson, J. Bishop, T. Sugimura, M. Terada, K. Toyoshima, and P. K. Vogt (ed).

30

-4-

- Japan Scientific Press, Tokyo). Microinjection into NIH 3T3 cells of oncogenically activated but not wild-type versions of the Raf-protein prepared with *Escherichia coli* expression vectors results in morphological transformation and stimulates DNA synthesis (Rapp, U.R., et al. (1987) In *Oncogenes and cancer*, S. A. Aaronson, J. Bishop, T. Sugimura, M. Terada, K. Toyoshima, and P. K. Vogt (ed.) Japan Scientific Press, Tokyo; Smith, M. R., et al (1990) *Mol. Cell. Biol.* 10:3828-3833). Activating mutants of B-Raf have been identified in a wide range of human cancers e.g. colon, ovarian, melanomas and sarcomas (Davies, H., et al. (2002), *Nature* 417 949-945. Published online June 9, 2002, 10.1038/nature00766). The preponderant mutation is a single phosphomimetic substitution in the kinase activation domain (V599E), leading to constitutive kinase activity and transformation of NIH3T3 cells.

- Thus, activated Raf-1 is an intracellular activator of cell growth. Raf-1 protein serine kinase in a candidate downstream effector of mitogen signal transduction, since Raf oncogenes overcome growth arrest resulting from a block of cellular ras activity due either to a cellular mutation (ras revertant cells) or microinjection of anti-ras antibodies (Rapp, U.R., et al. (1988) In *The Oncogene Handbook*, T. Curran, E.P. Reddy, and A. Skalka (ed.), Elsevier Science Publishers, The Netherlands, pp. 213-253; Smith, M.R., et al. (1986) *Nature* (London) 320:540-543).

- c-Raf function is required for transformation by a variety of membrane-bound oncogenes and for growth stimulation by mitogens contained in serums (Smith, M.R., et al. (1986) *Nature* (London) 320:540-543). Raf-1 protein serine kinase activity is regulated by mitogens via phosphorylation (Morrison, D.K., et al. (1989) *Cell* 58:648-657), which also effects subcellular distribution (Olah, Z., et al. (1991) *Exp. Brain Res.* 84:403; Rapp, U.R., et al. (1988) *Cold Spring Harbor Sym. Quant. Biol.* 53:173-184. Raf-1 activating growth factors include platelet-derived growth factor (PDGF) (Morrison, D.K., et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8855-8859),

-5-

- colony-stimulating factor (Baccarini, M., et al. (1990) *EMBO J.* 9:3649-3657), insulin (Blackshear, P.J., et al. (1990) *J. Biol. Chem.* 265:12115-12118), epidermal growth factor (EGF) (Morrison, R.K., et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8855-8859), interleukin 2 (Turner, B.C., et al (1991) *Proc. Natl. Acad. Sci. USA* 88:1227), and interleukin 3 and granulocytemacrophage colony-stimulating factor (Carroll, M.P., et al (1990) *J. Biol. Chem.* 265:19812-19817).

- Upon mitogen treatment of cells, the transiently activated Raf-1 protein serine kinase translocates to the perinuclear area and the nucleus (Olah, Z., et al. (1991) *Exp. Brain Res.* 84:403; Rapp, U.R., et al. (1988) *Cold Spring Harbor Sym. Quant. Biol.* 53:173-184). Cells containing activated Raf are altered in their pattern of gene expression (Heidecker, G., et al. (1989) In *Genes and signal transduction in multistage carcinogenesis*, N. Colburn (ed.), Marcel Dekker, Inc., New York, pp. 339-374), and Raf oncogenes activate transcription from Ap-1/P/E α 3-dependent promoters in transient transfection assays (Jamal, S., et al (1990) *Science* 344:463-466; Kalbuchi, K., et al (1989) *J. Biol. Chem.* 264:20855-20858; Waslyk, C., et al. (1989) *Mol. Cell. Biol.* 9:2247-2250).

- There are at least two independent pathways for Raf-1 activation by extracellular mitogens: one involving protein kinase C (KC) and a second initiated by protein tyrosine kinases (Blackshear, P.J., et al. (1990) *J. Biol. Chem.* 265:12131-12134; Kovach, K.S., et al (1990) *J. Biol. Chem.* 265:12115-12118; Morrison, D.K., et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:8855-8859; Siegel, J.N., et al (1990) *J. Biol. Chem.* 265:18472-18480; Turner, B.C., et al (1991) *Proc. Natl. Acad. Sci. USA* 88:1227). In either case, activation involves Raf-1 protein phosphorylation. Raf-1 phosphorylation may be a consequence of a kinase cascade amplified by autophosphorylation or may be caused entirely by autophosphorylation initiated by binding of a putative activating ligand to the Raf-1 regulatory

domain, analogous to PKC activation by diacylglycerol (Nishizuka, Y. (1986) Science 233:305-312).

The process of angiogenesis is the development of new blood vessels, generally capillaries, from pre-existing vasculature. Angiogenesis is defined as involving (i) activation of endothelial cells; (ii) increased vascular permeability; (iii) subsequent dissolution of the basement membrane and extravasation of plasma components leading to formation of a provisional fibrin gel extracellular matrix; (iv) proliferation and mobilization of endothelial cells; (v) reorganization of mobilized endothelial cells to form functional capillaries; (vi) capillary loop formation; and (vii) deposition of basement membrane and recruitment of perivascular cells to newly formed vessels.

Normal angiogenesis is activated during tissue growth, from embryonic development through maturity, and then enters a period of relative quiescence during adulthood.

Normal angiogenesis is also activated during wound healing, and at certain stages of the female reproductive cycle. Inappropriate or pathological angiogenesis has been associated with several disease states including various retinopathies; ischemic disease; atherosclerosis; chronic inflammatory disorders; rheumatoid arthritis, and cancer. The role of angiogenesis in disease states is discussed, for instance, in Fan et al, Trends in Pharmacol Sci. 16:54 66; Shawver et al, DOT Vol. 2, No. 2 February 1997; Folkmann, 1995, Nature Medicine 1:27-31.

In cancer the growth of solid tumors has been shown to be angiogenesis dependent. (See Folkmann, J., J. Natl. Cancer Inst., 1990, 82, 4-6.) Consequently, the targeting of pro-angiogenic pathways is a strategy being widely pursued in order to provide new therapeutics in these areas of great, unmet medical need.

Raf is involved in angiogenic processes. Endothelial growth factors (e.g. vascular endothelial growth factor VEGF or basic fibroblast growth factor bFGF) activates receptor tyrosine kinases (e.g. VEGFR-2) and signal through the Ras/Raf/Mek/Erk kinase cascade and protects endothelial cells from apoptosis (Alavi et al. (2003), Science 301, 94-96; Hood, J.D. et al. (2002), Science 296, 2404; Mikula, M. et al. (2001), EMBO J. 20, 1952; Hauser, M. et al. (2001), EMBO J. 20, 1940; Wojnowski et al. (1997), Nature Genet. 16, 293). Activation of VEGFR-2 by VEGF is a critical step in the signal transduction pathway that initiates tumor angiogenesis. VEGF expression may be constitutive to tumor cells and can also be upregulated in response to certain stimuli. One such stimuli is hypoxia, where VEGF expression is upregulated in both tumor and associated host tissues. The VEGF ligand activates VEGFR-2 by binding with its extracellular VEGF binding site. This leads to receptor dimerization of VEGFRs and autophosphorylation of tyrosine residues at the intracellular kinase domain of VEGFR-2. The kinase domain operates to transfer a phosphate from ATP to the tyrosine residues, thus providing binding sites for signaling proteins downstream of VEGFR-2 leading ultimately to initiation of angiogenesis (McMahon, G., The Oncologist, Vol. 5, No. 90001, 3-10, April 2000).

Mice with a targeted disruption in the Braf gene die of vascular defects during development (Wojnowski, L. et al. 1997, Nature genetics 16, page 293- 296). These mice show defects in the formation of the vascular system and in angiogenesis e.g. enlarged blood vessels and increased apoptotic death of differentiated endothelial cells.

For the identification of a signal transduction pathway and the detection of cross talks with other signaling pathways suitable models or model systems have been generated by various scientists, for example cell culture models (e.g. Khwaja et al., EMBO, 1997, 16, 2783-93) and

- 8 -

transgenic animal models (e.g. White et al., *Oncogene*, 2001, 20, 7064-7072). For the examination of particular steps in the signal transduction cascade, interfering compounds can be used for signal modulation (e.g. Stephens et al., *Biochemical J.*, 2000, 351, 95-105). The compounds according to the invention may also be useful as reagents for the examination of kinase dependent signal transduction pathways in animal and/or cell culture models or any of the clinical disorders listed throughout this application.

10 The measurement of kinase activity is a well known technique feasible for each person skilled in the art. Generic test systems for kinase activity detection with substrates, for example histone (e.g. Alessi et al., *FEBS Lett.* 1996, 399, 3, page 333-8) or myelin basic protein are well described in the literature (e.g. Campos-González, R. and Glenney, Jr., *J.R.* 1992 *J. Biol. Chem.* 267, Page 14535).

20 For the identification of kinase inhibitors various assay systems are available (see for example Walters et al., *Nature Drug Discovery* 2003, 2, page 259-266). For example, in scintillation proximity assays (e.g. Sorg et al., *J. of Biomolecular Screening*, 2002, 7, 11-19) or flashplate assays the radioactive phosphorylation of a protein or peptide as substrate with γ -ATP can be measured. In the presence of an inhibitory compound no signal or a decreased radioactive signal is detectable. Furthermore homogeneous time-resolved fluorescence resonance energy transfer (HTR-FRET), and fluorescence polarization (FP) technologies are useful for assay methods (for example Silis et al., *J. of Biomolecular Screening*, 2002, 191-214).

30 Other non-radioactive ELISA based assay methods use specific phospho-antibodies (AB). The phospho-AB binds only the phosphorylated substrate. This binding is detectable with a secondary peroxidase conjugated

- 9 -

antibody, measured for example by chemiluminescence (for example Ross et al., *Biochem. J.*, 2002, 366, 977-981).

5 The present invention provides compounds generally described as methylene urea derivatives, including both aryl and/or heteroaryl derivatives which are preferably kinase inhibitors and more preferably inhibitors of the enzyme raf kinase. Since the enzyme is a downstream effector of p21^{ras}, the inhibitors are useful in pharmaceutical compositions for human or veterinary use where inhibition of the raf kinase pathway is indicated, e.g., in the treatment of tumors and/or cancerous cell growth mediated by raf kinase. In particular, the compounds are useful in the treatment of human or animal solid cancers, e.g. murine cancer, since the progression of these cancers is dependent upon the ras protein signal transduction cascade and therefore susceptible to treatment by interruption of the cascade, i.e., by inhibiting raf kinase. Accordingly, the compound of Formula I or a pharmaceutically acceptable salt thereof is administered for the treatment of diseases mediated by the raf kinase pathway especially cancers, including solid cancers, such as, for example, carcinomas (e.g., of the lungs, pancreas, thyroid, bladder or colon), myeloid disorders (e.g., myeloid leukemia) or adenomas (e.g., villous colon adenoma), pathological angiogenesis and metastatic cell migration. Furthermore the compounds are useful in the treatment of complement activation dependent chronic inflammation (Niculescu et al. (2002) *Immunol. Res.*, 24:191-199) and HIV-1 (human immunodeficiency virus type1) induced immunodeficiency (Popik et al. (1998) *J. Virol.* 72: 6406-6413) and infection disease, Influenza A virus (Pleschka, S. et al. (2001), *Nat. Cell. Biol.*, 3(3):301-5) and *Helicobacter pylori* infection (Messler, S. et al. (2002), *FASEB J.*, 16(3): 417-9).

30 Therefore, subject of the present invention are methylene urea derivatives of formula I

group consisting of nitrogen, oxygen and sulfur, wherein L' is preferably substituted by at least one substituent selected from the group consisting of $-SO_2R_x$, $-C(O)R_x$ and $-C(NR_y)R_z$,

5 B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms, preferably of up to 20 carbon atoms, comprising at least one 5-, 6-, or 7-membered cyclic structure, preferably a 5- or 6-membered cyclic structure, bound directly to D containing 0-4 members of the group consisting of nitrogen, oxygen and sulfur, wherein said cyclic structure directly bound to D is preferably selected from the group consisting of aryl, heteroaryl and heterocyclyl, R_y is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally halosubstituted, up to per halo,

15 R_z is hydrogen or a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen;

R_x is R_z or NR_yR_z , where R_a and R_b are

25 a) independently hydrogen, a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms, selected from N, S and O, and are optionally substituted by halogen, or

30

A-D-B (I)

wherein

5 D is a bivalent methylene urea moiety which is directly bonded to A and B, preferably to one bonding partner via the carbon atom of the N-methylene moiety and to the other bonding partner via the N'-nitrogen atom, wherein the carbon atom of the N-methylene moiety is unsubstituted or substituted with one or more substituents, wherein said substituents are preferably selected from the group consisting of alkyl, alkylene, halogen, haloalkyl, C_3-C_7 -cycloalkyl, C_3-C_7 -cycloalkylene, heterocyclyl, aryl, aralkyl, heteroaryl, hydroxy, alkoxy, haloalkoxy, aralkoxy, aryloxy, mercapto, alkylsulfanyl, haloalkylsulfanyl, arylsulfanyl, heteroaryl/sulfanyl, alkylsulfenyl, haloalkylsulfenyl, arylsulfenyl, heteroaryl/sulfenyl, alkylsulfonfyl, haloalkylsulfonfyl, arylsulfonfyl, heteroaryl/sulfonfyl, carboxy, cyano, cyanoalkyl, aminosulfonfyl, acyl, acyloxy, carbamoyl, aryl, heteroaryl, heteroaryloxy, unsubstituted amino groups and substituted amino groups, and wherein the carbonyl group of said methylene urea moiety can be derivatized, preferably to a $C=S$, $C=NR^5$, $C=C(R^5)-NO_2$, $C=C(R^5)-CN$ or $C=C(CN)_2$ group

A is a unsubstituted or preferably substituted moiety of up to 40 carbon atoms of the formula: $-L-(M-L')_n$, where L is a 5, 6 or 7 membered cyclic structure, preferably selected from the group consisting of aryl, heteroaryl, arylene and heteroarylene, bound directly to D, L' comprises an optionally substituted cyclic moiety having at least 5 members, preferably selected from the group consisting of aryl, heteroaryl, aralkyl, cycloalkyl and heterocyclyl, M is a bond or a bridging group having at least to one atom, α is an integer of from 1-4; and each cyclic structure of L and L' contains 0-4 members of the

30

- 12 -

5 -OSi(R)₃ where R_i is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O, and are optionally substituted by halogen; or

10 b) R_a and R_b together from a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, or a substituted 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen; or

15 c) one of R_a or R_b is -C(O)-, a C₁-C₆ divalent alkylene group or a substituted C₁-C₆ divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted C₁-C₆ divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen; where B is substituted, L is substituted or L' is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo and W₁, where γ is 0-3;

25 wherein each W is independently selected from the group consisting of -CN, -CO₂R, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵, -Q-Ar, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O and optionally

30

- 13 -

substituted by one or more substituents independently selected from the group consisting of -CN, -CO₂R, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵ and halogen up to per-halo; with each R⁵ independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S

5

10 d) and O and optionally substituted by halogen; wherein Q is -O-, -S-, -N(R⁵)-, -(CH₂)_β-, -C(O)-, -CH(OH)-, -(CH₂)_β-, -(CH₂)_βS-, -(CH₂)_βN(R⁵)-, -O(CH₂)_β-CHHal-, -CHalZ-, -S-(CH₂)_β and -N(R⁵)(CH₂)_β where $\beta = 1-3$, and Hal is halogen; and

15 Ar is a 5- or 6-member aromatic structure containing 0-2 members selected from the group consisting of nitrogen, oxygen and sulfur, which is optionally substituted by halogen, up to per-halo, and optionally substituted by Z₁; wherein $\delta 1$ is 0 to 3 and each Z is independently selected from the group consisting of -CN, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, A and O and optionally substituted by one or more substituents selected from the group consisting of -CN, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵, and with R⁵ as defined above,

25 and the pharmaceutically acceptable derivatives, solvates, salts and stereoisomers thereof, including mixtures thereof in all ratios, and more preferred the salts and/or solvates thereof, and especially preferred the physiologically acceptable salts and/or solvates thereof.

30

More preferred, in the compound of formula I,

R_y is hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, C_{3-10} cycloalkyl having 0-3 heteroatoms, C_{2-10} alkenyl, C_{1-10} alkenoyl, C_{6-12} aryl, C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, C_{7-24} aralkyl, substituted C_{1-10} alkyl, substituted C_{1-10} alkoxy, substituted C_{3-10} cycloalkyl having 0-3 heteroatoms selected from N, S and O, substituted C_{6-14} aryl, substituted C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, substituted C_{7-24} aralkyl or substituted C_{7-24} aralkyl, where R_y is a substituted group, it is substituted by halogen up to per halo,

R_z is hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, C_{3-10} cycloalkyl having 0-3 heteroatoms, C_{2-10} alkenyl, C_{1-10} alkenoyl, C_{6-12} aryl, C_{3-12} hetaryl having 1-3 heteroatoms selected from S, N and O, C_{7-24} aralkyl, C_{7-24} aralkyl, substituted C_{3-10} cycloalkyl having 0-3 heteroatoms selected from S, N and O, substituted C_{3-12} hetaryl having 1-3 heteroatoms selected from S, N and O, substituted C_{7-24} aralkyl or substituted C_{7-24} aralkyl, where R_z is a substituted group, it is substituted by halogen up to per halo, hydroxy, C_{1-10} alkyl, C_{3-12} cycloalkyl having 0-3 heteroatoms selected from N, S and O, substituted C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, C_{7-24} aralkyl, C_{6-12} halo substituted alkyl up to per halo alkyl, C_{6-12} halo substituted aryl up to per halo aryl, C_{3-12} halo substituted cycloalkyl up to per halo cycloalkyl having 0-3 heteroatoms selected from N, S and O, halo substituted C_{3-12} hetaryl up to per halo, hetaryl having 1-3 heteroatoms selected from O, N and S, halo substituted C_{7-24} aralkyl up to per halo aralkyl, halo substituted C_{7-24} aralkyl up to per halo alkyl, and $-C(O)R_9$,

R_a and R_b are,

a) independently hydrogen, a carbon based moiety selected from the group consisting of C_{1-10} alkyl, C_{1-10} alkoxy, C_{3-10} cycloalkyl, C_{2-10} alkenyl, C_{1-10} alkenoyl, C_{6-12} aryl, C_{3-12} hetaryl having 1-3 heteroatoms selected from O, N and S, C_{3-12} cycloalkyl having 0-3 heteroatoms selected from N, S and O, C_{7-24} aralkyl, C_{7-24} aralkyl, substituted C_{1-10} alkyl, substituted C_{1-10} alkoxy, substituted C_{3-10} cycloalkyl, having 0-3 heteroatoms selected from N, S and O, substituted C_{6-12} aryl, substituted C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, substituted C_{7-24} aralkyl, where R_a and R_b are a substituted group, they are substituted by halogen up to per halo, hydroxy, C_{1-10} alkyl, C_{3-12} cycloalkyl having 0-3 heteroatoms selected from O, S and N, C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, C_{1-10} alkoxy, C_{6-12} aryl, C_{1-6} halo substituted alkyl up to per halo alkyl, C_{6-12} halo substituted aryl up to per halo aryl, C_{3-12} halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted C_{3-12} hetaryl up to per halo heteraryl, halo substituted C_{7-24} aralkyl up to per halo aralkyl, halo substituted C_{7-24} aralkyl up to per halo alkyl, and $-C(O)R_9$, or $-OSi(R_1)_3$ where R_1 is hydrogen, C_{1-10} alkyl, C_{1-10} alkoxy, C_{3-10} cycloalkyl, C_{2-10} alkenyl, C_{1-10} alkenoyl, C_{6-12} aryl, C_{3-12} hetaryl having 1-3 heteroatoms selected from O, N and S, C_{3-12} cycloalkyl having 0-3 heteroatoms selected from N, S and O, C_{7-24} aralkyl, C_{7-24} aralkyl, substituted C_{1-10} alkyl, substituted C_{1-10} alkoxy, substituted C_{3-10} cycloalkyl, having 0-3 heteroatoms selected from N, S and O, substituted C_{6-12} aryl, substituted C_{3-12} hetaryl having 1-3 heteroatoms selected from N, S and O, substituted C_{7-24} aralkyl, where R_a and R_b are a substituted group, they are substituted by halogen up to per halo, hydroxy, C_{1-10} alkyl, C_{3-12} cycloalkyl having 0-3

- 16 -

heteroatoms selected from O, S and N, C₃₋₁₂ hetaryl having 1-3 heteroatoms selected from N, S and O, C₁₋₁₀ alkoxy, C₆₋₁₂ aryl, C₁₋₆ halo substituted alkyl up to per halo alkyl, C_{6-C12} halo substituted aryl up to per halo aryl, C_{3-C12} halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted C_{3-C12} hetaryl up to per halo heteraryl, halo substituted C_{7-C24} aralkyl up to per halo aralkyl, halo substituted C_{7-C24} alkaryl up to per halo alkaryl, and -C(O)R₉.

or

b) R₈ and R₉ together form a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, or a substituted 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O with substituents selected from the group consisting of halogen up to per halo, hydroxy, C_{1-C10} alkyl, C_{1-C10} alkoxy, C₃₋₁₀ cycloalkyl, C₂₋₁₀ alkenyl, C₁₋₁₀ alkenoyl, C₆₋₁₂ aryl, C₃₋₁₂ hetaryl having 1-3 heteroatoms selected from O, N and S, C₃₋₁₂ cycloalkyl having 0-3 heteroatoms selected from N, S and O, C₇₋₂₄ aralkyl, C_{7-C24} alkaryl, substituted C₁₋₁₀ alkyl, substituted C₁₋₁₀ alkoxy, substituted C₃₋₁₀ cycloalkyl, having 0-3 heteroatoms selected from N, S and O, substituted C₆₋₁₂ aryl, substituted C₃₋₁₂ hetaryl having 1-3 heteroatoms selected from N, S and O, substituted C₇₋₂₄ aralkyl, substituted C₇₋₂₄ alkaryl, where R₈ and R₉ are a substituted group, they are substituted by halogen up

c) to per halo, hydroxy, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl having 0-3 heteroatoms selected from O, S and N, C₃₋₁₂ hetaryl having 1-3 heteroatoms selected from N, S and O, C₁₋₁₀ alkoxy, C₆₋₁₂ aryl, C₁₋₆ halo substituted alkyl up to per halo alkyl, C_{6-C12} halo substituted aryl up to per halo aryl, C_{3-C12} halo substituted

- 17 -

cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted C_{3-C12} hetaryl up to per halo heteraryl, halo substituted C_{7-C24} aralkyl up to per halo aralkyl, halo substituted C_{7-C24} alkaryl up to per halo alkaryl, and -C(O)R₉,

or

d) one of R₈ or R₉ is -C(O)-, a C_{1-C5} divalent alkylene group or a substituted C_{1-C5} divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted C_{1-C5} divalent alkylene group are selected from the group consisting of halogen, hydroxy, C₁₋₁₀ alkyl, C₃₋₁₂ cycloalkyl having 0-3 heteroatoms selected from, S and N, C₃₋₁₂ hetaryl having 1-3 heteroatoms selected from N, S and O, C₁₋₁₀ alkoxy, C₆₋₁₂ aryl, C_{7-C24} alkaryl, C_{7-C24} aralkyl, C₁₋₆ halo substituted alkyl up to per halo alkyl, C_{6-C12} halo substituted aryl up to per halo aryl, C_{3-C12} halo substituted cycloalkyl having 0-3 heteroatoms selected from N, S and O, up to per halo cycloalkyl, halo substituted C_{3-C12} hetaryl up to per halo heteraryl, halo substituted C_{7-C24} aralkyl up to per halo aralkyl, halo substituted C_{7-C24} alkaryl up to per halo alkaryl, and -C(O)R₉.

where R₉ is C₁₋₁₀ alkyl, -CN, -CO₂R₈, -OR₈, -SR₈, -NO₂, -C(O)R₈, -NR₈R₉, -NR₈C(O)OR₈ and -NR₈(CO)R₈ and R₈ and R₉ are independently selected from the group consisting of hydrogen, C₁₋₁₀ alkyl, C₁₋₁₀ alkoxy, C₃₋₁₀ cycloalkyl having 0-3 heteroatoms selected from O, N and S, C₆₋₁₂ aryl, C_{3-C12} hetaryl with 1-3 heteroatoms selected from O, N and S and C_{7-C24} aralkyl, C_{7-C24} alkaryl, up to per halo substituted C_{1-C10} alkyl, up to per halo substituted C_{3-C10} cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per halo substituted C_{6-C14} aryl, up to per halo substituted C_{3-C12} hetaryl having 1-3 heteroatoms

selected from O, N and S, halo substituted C₇-C₂₄ alkaryl up to per halo alkaryl, and up to per halo substituted C₇-C₂₄ alkaryl,

W is independently selected from the group consisting -CN, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, C₂-C₁₀ alkenyl, C₁-C₁₀ alkenyl, C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, S and N, C₆-C₁₄ aryl, C₇-C₂₄ alkaryl, C₇-C₂₄ alkaryl, C₃-C₁₂ heteroaryl having 1-3 heteroatoms selected from O, N and S, C₄-C₂₃ alkheteroaryl having 1-3 heteroatoms selected from O, N and S, substituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkoxy, substituted C₂-C₁₀ alkenyl, substituted C₁-C₁₀ alkenyl, substituted C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, N and S, substituted C₇-C₂₄ alkaryl, substituted C₇-C₂₄ alkaryl, substituted C₄-C₂₃ alkheteroaryl having 1-3 heteroatoms selected from O, N and S, and -Q-Ar,

R⁵ is independently selected from H, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, C₂-C₁₀ alkenyl, C₁-C₁₀ alkenyl, C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, C and N, C₆-C₁₄ aryl, C₃-C₁₃ hetaryl having 1-3 heteroatoms selected from O, N and S, C₇-C₁₄ alkaryl, C₇-C₂₄ alkaryl, C₄-C₂₃ alkheteroaryl having 1-3 heteroatoms selected from O, N and S, up to per-halosubstituted C₁-C₁₀ alkyl, up to per-halosubstituted C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, N and S, up to per-halosubstituted C₆-C₁₄ aryl, up to per-halosubstituted C₃-C₁₃ hetaryl having 1-3 heteroatoms selected from O, N and S, up to per-halosubstituted C₇-C₂₄ alkaryl, up to per-halosubstituted C₄-C₂₃ alkheteroaryl; and each

Z is independently selected from the group consisting -CN, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵,

-NR⁵C(O)R⁵, C₁-C₁₀ alkyl, C₁-C₁₀ alkoxy, C₂-C₁₀ alkenyl, C₁-C₁₀ alkenyl, C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, S and N, C₆-C₁₄ aryl, C₇-C₂₄ alkaryl, C₇-C₂₄ alkaryl, C₃-C₁₂ heteroaryl having 1-3 heteroatoms selected from O, N and S, C₄-C₂₃ alkheteroaryl having 1-3 heteroatoms selected from O, N and S, substituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkoxy, substituted C₂-C₁₀ alkenyl, substituted C₁-C₁₀ alkenyl, substituted C₃-C₁₀ cycloalkyl having 0-3 heteroatoms selected from O, N and S, substituted C₆-C₁₂ aryl, substituted C₃-C₁₂ hetaryl having 1-3 heteroatoms selected from O, N and S; wherein if Z is a substituted group, the one or more substituents are selected from the group consisting of -CN, -CO₂R⁵, -C(O)NR⁵R⁵, -C(O)R⁵, -NO₂, -OR⁵, -SR⁵, -NR⁵R⁵, -NR⁵C(O)OR⁵, -NR⁵C(O)R⁵.

According to the invention, each M independently from one another represents a bond OR is a bridging group, selected from the group consisting of (CR⁵R⁵)_n, or (CHR⁵)_n-Q-(CHR⁵)_n, wherein

Q is selected from a group consisting of O, S, N-R⁵, (CHal₂)_n, (O-CHR⁵)_n, (CHR⁵-O)_n, CR⁵=CR⁵, (O-CHR⁵CHR⁵)_n, (CHR⁵CHR⁵-O)_n, C=O, C=S, C=NR⁵, CH(OR⁵), C(OR⁵)(OR⁵), C(=O)O, OC(=O), OC(=O)O, C(=O)N(R⁵), N(R⁵)C(=O), OC(=O)N(R⁵), N(R⁵)C(=O)O, CH=N-O, CH=N-NR⁵, OC(O)NR⁵, NR⁵C(O)O, S=O, SO₂, SO₂NR⁵ and NR⁵SO₂, wherein

R⁵ is in each case independently selected from the meanings given above, preferably from hydrogen, halogen, alkyl, aryl, alkaryl,

h, i are independently from each other 0, 1, 2, 3, 4, 5 or 6, preferably 0, 1, 2, or 3, and

j is 1, 2, 3, 4, 5 or 6, preferably 0, 1, 2 or 3.

- 20 -

- More preferred, each M independently from one another represents a bond or is a bridging group, selected from the group consisting of $-O-$, $-S-$, $-N(R^5)-$, $-(CH_2)_\beta-$, $-C(O)-$, $-CH(OH)-$, $-(CH_2)_\beta O-$, $-(CH_2)_\beta S-$, $-(CH_2)_\beta N(R^5)-$, $-O(CH_2)_\beta$, $-CHHal-$, $-CHaI_2-$, $-S-(CH_2)_\beta-$ and $-N(R^5)(CH_2)_\beta$, where β is 1 to 6 and especially preferred 1 to 3, Hal is halogen and R^5 is as defined above.
- More preferred, the group B of Formula I is a substituted or unsubstituted six member aryl moiety or six member heteraryl moiety, said heteraryl moiety having 1 to 4 members selected from the group of heteraryl atoms consisting of nitrogen, oxygen and sulfur with the balance of the heteraryl moiety being carbon.

Even more preferred, the group B of Formula I is

- 15 a) an unsubstituted phenyl group, an unsubstituted pyridyl group, an unsubstituted pyrimidinyl, a phenyl group substituted by a substituent selected from the group consisting of halogen and W_γ wherein W and γ are as defined in claim 1, a pyrimidinyl group substituted by a substituent selected from the group constituting of halogen and W_γ , whereas W and γ are as defined above, or a substituted pyridyl group, substituted by a substituent selected from the group consisting of halogen and W_γ wherein W and γ are as defined above; or a substituted phenyl group, a substituted pyrimidinyl group, or substituted pyridyl group substituted 1 to 3 times by 1 or more substituents selected from the group consisting of $-CN$, halogen, C_1-C_{10} alkyl, C_1-C_{10} alkyl alkoxy, $-OH$, up to per halo substituted C_1-C_{10} alkyl, up to per halo substituted C_1-C_{10} alkoxy or phenyl substituted by halogen up to per halo; or
- 20 b) a substituted phenyl group, a substituted pyrimidinyl group, or substituted pyridyl group substituted 1 to 3 times b 1 or more

- 21 -

- substituents selected from the group consisting of $-CN$, halogen, alkyl, especially C_1-C_4 alkyl, alkoxy, especially C_1-C_4 alkoxy, $-OH$, up to per halo substituted alkyl, especially up to per halo substituted C_1-C_4 alkyl, up to per halo substituted alkoxy, especially up to per halo substituted C_1-C_4 alkoxy or phenyl substituted by halogen up to per halo.

- 10 In the formula I, the group L which is directly bound to D is preferably a substituted or unsubstituted 6 member aryl moiety or a substituted or unsubstituted 6 member heteraryl moiety, wherein said heteraryl moiety has 1 to 4 members selected from the group of heteroatoms consisting of nitrogen, oxygen and sulfur with the balance of said heteraryl moiety being carbon, wherein the one or more substituents are selected from the group consisting of halogen and W_γ wherein W and γ are as defined above.

- 15 More preferred, the group L is a substituted phenyl, unsubstituted phenyl, substituted pyrimidinyl, unsubstituted pyrimidinyl, substituted pyridyl or unsubstituted pyridyl group.

- 20 In the formula I, the group L' preferably comprises a 5 to 6 membered aryl moiety or heteraryl moiety, wherein said heteraryl moiety comprises 1 to 4 members selected from the group of heteroatoms consisting of nitrogen, oxygen and sulfur.

- 25 More preferred, the group L' is phenyl, pyridinyl or pyrimidinyl.

- 30 According to the invention, a methylene moiety is a bivalent radical of formula $-CRR_2-$, where R and R_2 are selected independently from one another from hydrogen or suitable substituents other than hydrogen. Suitable substituents are preferably selected from the group consisting of alkyl, alkylene, halogen, haloalkyl, C_3-C_7 -cycloalkyl, C_3-C_7 -cycloalkylene,

wherein A and B are as defined above/below, and wherein the carbonyl moiety in formula Ia can be derivatized as described above/below, and the salts or solvates thereof. Especially preferred are compounds of formula Ia, wherein the carbonyl moiety is not derivatized.

Preferably, A or B is substituted by one or more substituents as described above/below. More preferably, A and B each are substituted by one or more substituents as described above/below. Even more preferably, A is substituted by two or more substituents as described above/below.

Preferably, subject of the present invention are the optically active forms or stereo isomers of the compounds according to the invention, such as the enantiomers, the diastereomers and/or mixtures thereof in all ratios, such as, for example, stereochemically uniform compounds or racemates. Preferably, further subject of the present invention are the solvates and hydrates of the compounds according to the invention. Preferably, further subject of the present invention are the pharmaceutically acceptable derivatives or physiologically functional derivatives of the compounds according to the invention. More preferably, further subject of the present invention are the salts of the compounds according to the invention, especially the pharmaceutically and/or physiologically acceptable salts of compounds according to the invention.


As used herein, the term "effective amount" means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue, system, animal or human that is being sought, for instance, by a researcher or clinician. Furthermore, the term "therapeutically effective amount" means any amount which, as compared to a corresponding subject who has not received such amount, results in improved treatment, healing, prevention, or amelioration of a disease, disorder, or side effect, or a decrease in the rate of advancement of a disease or disorder. The

heterocyclyl, aryl, aralkyl, heteroaryl, hydroxy, alkoxy, haloalkoxy, aralkoxy, aryloxy, mercapto, alkylsulfanyl, haloalkylsulfanyl, arylsulfanyl, heteroarylsulfanyl, alkylsulfenyl, haloalkylsulfenyl, arylsulfenyl, heteroarylsulfenyl, alkylsulfonyl, haloalkylsulfonyl, arylsulfonyl, heteroarylsulfonyl, carboxy, cyano, cyanoalkyl, aminosulfonyl, acyl, acyloxy, carbamoyl, aryl, heteroaryl, heteroaryloxy, unsubstituted amino groups and substituted amino groups. A methylene moiety wherein R and R both are hydrogen is also referred to as an unsubstituted methylene moiety. A methylene moiety wherein R and/or R are other than hydrogen is referred to as a substituted methylene moiety.

Thus, a methylene urea moiety according to the invention is a bivalent radical wherein one of the nitrogen atoms of the urea moiety is substituted by a methylene moiety. Preferably, A and B are bonded to the resulting methylene urea moiety via the nitrogen atom of the urea moiety that is not substituted by the methylene moiety, and to the carbon atom of the methylene moiety, respectively.

The hydrogen atoms of one or both nitrogen atoms of the methylene urea moiety can be substituted by suitable substituents, preferably selected from the group consisting of alkyl, alkylene, haloalkyl, C₃-C₇-cycloalkyl, C₃-C₇-cycloalkylene, heterocyclyl, aryl, aralkyl, heteroaryl, carboxy, cyanoalkyl, acyl and heteroaryl. Preferably, both nitrogen atoms of the methylene urea moiety are unsubstituted. In this respect, one or both of the nitrogen atoms of D can, independently from one another, optionally be deprotonated, protonated and/or quarternized. The resulting ions or salts are also subject of the present invention.

Accordingly, preferred compounds of formula I are of formula Ia



- 24 -

term also includes within its scope amounts effective to enhance normal physiological function.

- 5 As used herein, the term "alkyl" preferably refers to a straight or branched chain hydrocarbon having from one to twelve carbon atoms, optionally substituted with substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylsulfanyl, C₁-C₆ alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, or C₁-C₆ perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "alkyl" as used herein include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, t-butyl, n-pentyl, isopentyl, and the like.

- 15 As used herein, the term "C₁-C₆ alkyl" preferably refers to an alkyl group as defined above containing at least 1, and at most 6, carbon atoms.
- Examples of branched or straight chained "C₁-C₆ alkyl" groups useful in the present invention include, but are not limited to, methyl, ethyl, n-propyl, isopropyl, isobutyl, n-butyl, t-butyl, n-pentyl and isopentyl.
- 20 As used herein, the term "alkylene" preferably refers to a straight or branched chain divalent hydrocarbon radical having from one to ten carbon atoms, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfonyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl, optionally substituted by alkyl, nitro, cyano, halogen and lower perfluoroalkyl, multiple degrees of substitution being allowed.
- 25 Examples of "alkylene" as used herein include, but are not limited to, methylene, ethylene, n-propylene, n-butylene and the like.

- 30 As used herein, the term "C₁-C₆ alkylene" preferably refers to an alkylene group, as defined above, which contains at least 1, and at most 6, carbon

- 25 -

atoms respectively. Examples of "C₁-C₆ alkylene" groups useful in the present invention include, but are not limited to, methylene, ethylene and n-Propylene.

- 5 As used herein, the term "halogen" or "hal" preferably refers to fluorine (F), chlorine (Cl), bromine (Br) or iodine (I).
- 10 As used herein, the term "C₁-C₆ haloalkyl" preferably refers to an alkyl group as defined above containing at least 1, and at most 6, carbon atoms substituted with at least one halogen, halogen being as defined herein. Examples of branched or straight chained "C₁-C₆ haloalkyl" groups useful in the present invention include, but are not limited to, methyl, ethyl, propyl, isopropyl, isobutyl and n-butyl substituted independently with one or more halogens, e.g., fluoro, chloro, bromo and iodo.

- 15 As used herein, the term "C₃-C₇ cycloalkyl" preferably refers to a non-aromatic cyclic hydrocarbon ring having from three to seven carbon atoms and which optionally includes a C₁-C₆ alkyl linker through which it may be attached. The C₁-C₆ alkyl group is as defined above. Exemplary "C₃-C₇ cycloalkyl" groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl. The term "cycloalkyl", as used herein preferably also includes saturated heterocyclic groups, which are preferably selected from the cycloalkyl-groups as defined above, wherein one or two carbon atoms are replaced by hetero atoms, selected from the group consisting of O, N and S.
- 25

- 30 As used herein, the term "C₃-C₇ cycloalkylene" preferably refers to a non-aromatic alicyclic divalent hydrocarbon radical having from three to seven carbon atoms, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfanyl, lower alkylsulfonyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl,

aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, lower perfluoroalkyl, multiple degrees of substitution being allowed. Examples of "cycloalkylene" as used herein include, but are not limited to, cyclopropyl-1,1-diyl, cyclopropyl-1,2-diyl, cyclobutyl-1,2-diyl, cyclopentyl-1,3-diyl, cyclohexyl-1,4-diyl, cycloheptyl-1,4-diyl, or cyclooctyl-1,5-diyl, and the like.

5

As used herein, the term "heterocyclic" or the term "heterocyclyl" preferably refers to a three to twelve-membered heterocyclic ring having one or more degrees of unsaturation containing one or more heteroatomic substituents selected from S, SO, SO₂, O or N, optionally substituted with substituents selected from the group consisting of C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkylsulfonyl, C₁-C₆ haloalkylsulfonyl, C₁-C₆ alkoxy, C₁-C₆ alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, or C₁-C₆ perfluoroalkyl, multiple degrees of substitution being allowed. Such a ring may be optionally fused to one or more other "heterocyclic" ring(s) or cycloalkyl ring(s). Examples of "heterocyclic" moieties include, but are not limited to, tetrahydrofuran, pyran, 1,4-dioxane, 1,3-dioxane, pyrrolidine, piperidine, morpholine, tetrahydrothiopyran, tetrahydrothiophene, and the like.

15

20

As used herein, the term "heterocyclene" preferably refers to a three to twelve-membered heterocyclic ring diradical having one or more degrees of unsaturation containing one or more heteroatoms selected from S, SO, SO₂, O or N, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfonyl, lower alkylsulfonyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, nitro, cyano, halogen, lower perfluoroalkyl, multiple degrees of substitution being allowed. Such a ring may be optionally fused to one or more benzene rings or to one or more of

30

another "heterocyclic" rings or cycloalkyl rings. Examples of "heterocyclene" include, but are not limited to, tetrahydrofuran-2,5-diyl, morpholine-2,3-diyl, pyran-2,4-diyl, 1,4-dioxane-2,3-diyl, 1,3-dioxane-2,4-diyl, piperidine-2,4-diyl, piperidine-1,4-diyl, pyrrolidine-1,3-diyl, morpholine-2,4-diyl, and the like.

5

As used herein, the term "aryl" preferably refers to an optionally substituted benzene ring or to an optionally substituted benzene ring system fused to one or more optionally substituted benzene rings to form, for example, anthracene, phenanthrene, or naphthalene ring systems. Exemplary optional substituents include C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylsulfonyl, C₁-C₆ alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaryl, acyloxy, aroyloxy, heteroaryloxy, alkoxy, carbonyl, nitro, cyano, halogen, C₁-C₆ perfluoroalkyl, heteroaryl, or aryl, multiple degrees of substitution being allowed. Examples of "aryl" groups include, but are not limited to Phenyl, 2-naphthyl, 1-naphthyl, biphenyl, as well as substituted derivatives thereof. As used herein, the term "arylene" preferably refers to a benzene ring diradical or to a benzene ring system diradical fused to one or more optionally substituted benzene rings, optionally substituted with substituents selected from the group which includes lower alkyl, lower alkoxy, lower alkylsulfonyl, lower alkylsulfonyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaryl, acyloxy, aroyloxy, heteroaryloxy, alkoxy, carbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl and aryl, multiple degrees of substitution being allowed. Examples of "arylene" include, but are not limited to benzene-1,4-diyl, naphthalene-1,8-diyl, anthracene-1,4-diyl, and the like.

10

15

20

25

30

- 28 -

As used herein, the term "aralkyl" preferably refers to an aryl or heteroaryl group, as defined herein, attached through a C₁-C₆ alkyl linker, wherein C₆ alkyl is as defined herein. Examples of "aralkyl" include, but are not limited to, benzyl, phenylpropyl, 2-pyridylmethyl, 3-isoxazolylmethyl, 5-methyl-3-isoxazolylmethyl and 2-imidazolylmethyl.

As used herein, the term "heteroaryl" preferably refers to a monocyclic five to seven-membered aromatic ring, or to a fused bicyclic aromatic ring system comprising two of such monocyclic five to seven-membered aromatic rings. These heteroaryl rings contain one or more nitrogen, sulfur and/or oxygen heteroatoms, where N-Oxides and sulfur Oxides and dioxides are permissible heteroatom substitutions and may be optionally substituted with up to three members selected from a group consisting of C₁-C₆ alkyl, C₁-C₆ haloalkyl, C₁-C₆ alkoxy, C₁-C₆ alkylsulfanyl, C₁-C₆ haloalkylsulfanyl, C₁-C₆ alkylsulfenyl, C₁-C₆ alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl optionally substituted by alkyl, acyl, aroyl, heteroaryl, acyloxy, aryloxy, heteroaryloxy, alkoxy, carbonyl, nitro, cyano, halogen, C₁-C₆ perfluoroalkyl, heteroaryl or aryl, multiple degrees of substitution being allowed. Examples of "heteroaryl" groups used herein include furanyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, oxo-pyridyl, thiadiazolyl, isothiazolyl, pyridyl, pyridazyl, pyrazinyl, pyrimidyl, quinoliny, isoquinoliny, benzofuranyl, benzothiophenyl, indolyl, indazolyl, and substituted versions thereof.

As used herein, the term "heteroarylene" preferably refers to a five- to seven-membered aromatic ring diradical, or to a polycyclic heterocyclic aromatic ring diradical, containing one or more nitrogen, oxygen, or sulfur heteroatoms, where N-Oxides and sulfur monoxides and sulfur dioxides are permissible heteroatomic substitutions, optionally substituted with substituents selected from the group consisting of lower alkyl, lower

- 29 -

alkoxy, lower alkylsulfanyl, lower alkylsulfenyl, lower alkylsulfonyl, oxo, hydroxy, mercapto, amino optionally substituted by alkyl, carboxy, tetrazolyl, carbamoyl optionally substituted by alkyl, aminosulfonyl, optionally substituted by alkyl, acyl, aroyl, heteroaryl, acyloxy, aryloxy, heteroaryloxy, alkoxy, carbonyl, nitro, cyano, halogen, lower perfluoroalkyl, heteroaryl, or aryl, multiple degrees of substitution being allowed. For polycyclic aromatic ring system diradicals, one or more of the rings may contain one or more heteroatoms. Examples of "heteroarylene" used herein are furan-2,5-diyl, thiophene-2,4-diyl, 1,3,4-oxadiazole-2,5-diyl, 1,3,4-thiadiazole-2,5-diyl, 1,3-thiazole-2,5-diyl, pyridine-2,4-diyl, pyridine-2,3-diyl, pyridine-2,5-diyl, pyrimidine-2,4-diyl, quinoline-2,3-diyl, and the like.

As used herein, the term "alkoxy" preferably refers to the group R_aO-, where R_a is alkyl as defined above and the term "C₁-C₆ alkoxy" preferably refers to an alkoxy group as defined herein wherein the alkyl moiety contains at least 1 and at most 6 carbon atoms. Exemplary C₁-C₆ alkoxy groups useful in the present invention include, but are not limited to methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy and t-butoxy.

As used herein, the term "haloalkoxy" preferably refers to the group R_aO-, where R_a is haloalkyl as defined above and the term "C₁-C₆ haloalkoxy" preferably refers to a haloalkoxy group as defined herein wherein the haloalkyl moiety contains at least 1 and at most 6 carbon atoms. Exemplary C₁-C₆ haloalkoxy groups useful in the present invention include, but are not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy and t-butoxy substituted with one or more halo groups, for instance trifluoromethoxy.

As used herein the term "aralkoxy" preferably refers to the group R_cR_aO-, where R_a is alkyl and R_c is aryl as defined above.

As used herein the term "aryloxy" preferably refers to the group $R_C O-$, where R_C is aryl as defined above.

5 As used herein, the term "alkylsulfanyl" preferably refers to the group $R_A S-$, where R_A is alkyl as defined above and the term " C_1-C_6 alkylsulfanyl" preferably refers to an alkylsulfanyl group as defined herein wherein the alkyl moiety contains at least 1 and at most 6 carbon atoms.

10 As used herein, the term "haloalkylsulfanyl" preferably refers to the group $R_B S-$, where R_B is haloalkyl as defined above and the term " C_1-C_6 haloalkylsulfanyl" preferably refers to a haloalkylsulfanyl group as defined herein wherein the alkyl moiety contains at least 1 and at most 6 carbon atoms.

15 As used herein, the term "alkylsulfenyl" preferably refers to the group $R_A S(O)-$, where R_A is alkyl as defined above and the term " C_1-C_6 alkylsulfenyl" preferably refers to an alkylsulfenyl group as defined herein wherein the alkyl moiety contains at least 1 and at most 6 carbon atoms.

20 As used herein, the term "alkylsulfonyl" preferably refers to the group $R_A SO_2-$, where R_A is alkyl as defined above and the term " C_1-C_6 alkylsulfonyl" preferably refers to an alkylsulfonyl group as defined herein wherein the alkyl moiety contains at least 1 and at most 6 carbon atoms.

25 As used herein, the term "oxo" preferably refers to the group $=O$.

As used herein, the term "mercapto" preferably refers to the group $-SH$.

As used herein, the term "carboxy" preferably refers to the group $-COOH$.

30 As used herein, the term "cyano" preferably refers to the group $-CN$.

As used herein, the term "cyanoalkyl" preferably refers to the group $-R_B CN$, wherein R_B is alkyl as defined above. Exemplary "cyanoalkyl" groups useful in the present invention include, but are not limited to, cyanomethyl, cyanoethyl and cyanoisopropyl.

5 As used herein, the term "aminosulfonyl" preferably refers to the group $-SO_2 NH_2$.

10 As used herein, the term "carbamoyl" preferably refers to the group $-C(O)NH_2$.

As used herein, the term "sulfanyl" shall refer to the group $-S-$.

15 As used herein, the term "sulfenyl" shall refer to the group $-S(O)-$.

As used herein, the term "sulfonyl" shall refer to the group $-S(O)_2-$ or $-SO_2-$.

20 As used herein, the term "acyl" preferably refers to the group $R_F C(O)-$, where R_F is alkyl, cycloalkyl or heterocyclyl as defined herein.

As used herein, the term "aroyl" preferably refers to the group $R_C C(O)-$, where R_C is aryl as defined herein.

25 As used herein, the term "heteroaryl" preferably refers to the group $R_E C(O)-$, where R_E is heteroaryl as defined herein.

As used herein, the term "alkoxycarbonyl" preferably refers to the group $R_A OC(O)-$, where R_A is alkyl as defined herein.

30 As used herein, the term "acyloxy" preferably refers to the group $R_F C(O)O-$, where R_F is alkyl, cycloalkyl, or heterocyclyl as defined herein.

- 32 -

As used herein, the term "aryloxy" preferably refers to the group $R_C C(O)O$, where R_C is aryl as defined herein.

5 As used herein, the term "heteroaryloxy" preferably refers to the group $R_C C(O)O$, where R_C is heteroaryl as defined herein.

10 As used herein, the term "carbonyl" or "carbonyl moiety" preferably refers to the group $C=O$.

15 As used herein, the term "thiocarbonyl" or "thiocarbonyl moiety" preferably refers to the group $C=S$.

20 As used herein, the term "amino", "amino group" or "imino moiety" preferably refers to the group $NR_G R_G$, wherein R_G and R_G are preferably selected, independently from one another, from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkenyl, cycloalkyl, alkylencycloalkyl, cyanoalkyl, aryl, aralkyl, heteroaryl, acyl and aroyl. If both R_G and R_G are hydrogen, $NR_G R_G$ is also referred to as "unsubstituted amino moiety" or "unsubstituted amino group". If R_G and/or R_G are other than hydrogen, $NR_G R_G$ is also referred to as "substituted amino moiety" or "substituted amino group".

25 As used herein, the term "imino" or "imino moiety" preferably refers to the group $C=NR_G$, wherein R_G is preferably selected from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkenyl, cycloalkyl, alkylencycloalkyl, cyanoalkyl, aryl, aralkyl, heteroaryl, acyl and aroyl. If R_G is hydrogen, $C=NR_G$ is also referred to as "unsubstituted imino moiety". If R_G is a residue other than hydrogen, $C=NR_G$ is also referred to as "substituted imino moiety".

30

- 33 -

5 As used herein, the term "ethene-1,1-diyl moiety" preferably refers to the group $C=CR_R R_L$, wherein R_R and R_L are preferably selected, independently from one another, from the group consisting of hydrogen, halogen, alkyl, haloalkyl, alkenyl, cycloalkyl, nitro, alkylencycloalkyl, cyanoalkyl, aryl, haloalkyl, heteroaryl, acyl and aroyl. If both hydrogen R_R and R_L are hydrogen, $C=CR_R R_L$ is also referred to as "unsubstituted ethene-1,1-diyl moiety". If one of R_R and R_L or both are a residue other than hydrogen, $C=CR_R R_L$ is also referred to as "substituted ethene-1,1-diyl moiety".

10 As used herein, the terms "group", "residue" and "radical" or "groups", "residues" and "radicals" are usually used as synonyms, respectively, as it is common practice in the art.

15 As used herein, the term "optionally" means that the subsequently described event(s) may or may not occur, and includes both event(s), which occur, and events that do not occur.

20 As used herein, the term "pharmaceutically acceptable derivative" preferably refers to any physiologically functional derivative of a compound of the present invention, for example, an ester or an amide, which upon administration to a mammal is capable of providing (directly or indirectly) a compound of the present invention or an active metabolite thereof. Such derivatives are clear to those skilled in the art, without undue experimentation, and with reference to the teaching of Burger's Medicinal Chemistry And Drug Discovery, 5th Edition, Vol 1: Principles and Practice, which is incorporated herein by reference to the extent that it teaches physiologically functional derivatives. Such derivatives preferably include so-called prodrug-compounds, for example compounds according to the invention that are derivatized with alkyl groups, acyl groups, sugars or peptides, such as oligopeptides, and that are easily degraded or metabolized to the active compounds according to the invention. Such derivatives preferably include biodegradable polymer derivatives of the

25

30

compounds according to the invention. Suitable polymers and methods for producing biodegradable polymeric derivatives are known in the art, for example from Int. J. Pharm. 115, 61-67 (1995).

5 As used herein, the term "solvent" preferably refers to a complex of variable stoichiometry formed by a solute (in this invention, a compound of formula I or formula II or a salt or physiologically functional derivative thereof) and a solvent. Such solvents for the purpose of the invention may not interfere with the biological activity of the solute. Examples of suitable solvents include, but are not limited to, water, methanol, ethanol and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of suitable pharmaceutically acceptable solvents include, without limitation, water, ethanol and acetic acid. Most preferably the solvent used is water. Examples for suitable solvents are the mono- or dihydrates or alcoholates of the compounds according to the invention.

15 As used herein, the term "substituted" preferably refers to substitution with the named substituent or substituents, multiple degrees of substitution being allowed unless otherwise stated.

20 Certain of the compounds described herein may contain one or more chiral atoms, or may otherwise be capable of existing as two or more stereoisomers, which are usually enantiomers and/or diastereomers.

25 Accordingly, the compounds of this invention include mixtures of stereoisomers, especially mixtures of enantiomers, as well as purified stereoisomers, especially purified enantiomers, or stereoisomerically enriched mixtures, especially enantiomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds represented by formulae I and II above as well as any wholly or partially equilibrated mixtures thereof. The present invention also covers the individual isomers of the compounds represented by the formulas above as mixtures with isomers thereof in which one or more chiral

Centers are inverted. Also, it is understood that all tautomers and mixtures of tautomers of the compounds of formulae (I) or (II) are included within the scope of the compounds of formulae (I) and (II) and preferably the formulae and subformulae corresponding thereto.

5 Racemates obtained can be resolved into the isomers mechanically or chemically by methods known per se. Diastereomers are preferably formed from the racemic mixture by reaction with an optically active resolving agent. Examples of suitable resolving agents are optically active acids, such as the D and L forms of tartaric acid, diacetyltartaric acid, dibenzoyltartaric acid, mandelic acid, malic acid, lactic acid or the various optically active camphorsulfonic acids, such as β -camphorsulfonic acid. Also advantageous is enantiomer resolution with the aid of a column filled with an optically active resolving agent (for example dinitrobenzophenylglycine); an example of a suitable eluent is a hexane/isopropanol/acetone/nitrile mixture.

20 The diastereomer resolution can also be carried out by standard purification processes, such as, for example, chromatography or fractional crystallization.


It is of course also possible to obtain optically active compounds of the formula I or II by the methods described above by using starting materials which are already optically active.

25 Unless indicated otherwise, it is to be understood that reference to compounds of formula I preferably includes the reference to the compounds of formula II. Unless indicated otherwise, it is to be understood that reference to the compounds of formula II preferably includes the reference to the sub formulae corresponding thereto, for example the sub formulae II.1 to II.20 and preferably formulae IIa to IIx. It is also understood

30

that the following embodiments, including uses and compositions, although recited with respect to formula I are preferably also applicable to formulae II, sub formulae II.1 to II.20 and preferably formulae IIa to IIx.

5 Especially preferred compounds according to the invention are compounds of formula II



wherein

are selected independently from one another from aromatic hydrocarbons containing 6 to 14 carbon atoms and ethylenical unsaturated or aromatic heterocyclic residues containing 3 to 10 carbon atoms and one or two heteroatoms, independently selected from N, O and S,

R^6, R^7
 R^8, R^9 and R^{10} ,
are independently selected from the meanings given for
or R^6 and R^7 together form a carbocyclic residue
comprising 3 to 7 carbon atoms or a heterocyclic
residue comprising 1, 2 or 3 hetero atoms, selected
from the group consisting of O, N and S, and 2 to 6
carbon atoms, said carbocyclic or heterocyclic residue
being unsubstituted or comprising 1, 2 or 3 substituents,
selected from the meanings given for R^8, R^9 and R^{10} .

E, G, M, Q and U are selected, independently from one another, from carbon atoms and nitrogen atoms, with the proviso that one or more of E, G, M, Q and U are carbon atoms and that X is bonded to a carbon atom,


5. R^8, R^9 and R^{10} are independently selected from a group consisting of

- are independently selected from a group consisting of H, A, and $(\text{CH}_2)_m\text{Ar}^6$, wherein
- is a 5- or 6-membered aromatic hydrocarbon which is optionally substituted by one or more substituents selected from a group consisting of methyl, ethyl, propyl, 2-propyl, tert-butyl, Hal, CN, OH, NH₂ and CF₃,
- are independently of one another 0, 1, 2, 3, 4, or 5,
- represents a bond or is $(\text{CR}^{11}\text{R}^{12})_h$, or $(\text{CHR}^{11})_h\text{-Q-}$ $(\text{CHR}^{12})_h$, wherein
- is selected from a group consisting of O, S, N-R¹⁵, $(\text{CHAl}_2)_h$, $(\text{O-CHR}^{18})_h$, $(\text{CHR}^{18}\text{-O})_h$, $\text{CR}^{18}=\text{CR}^{19}$, $(\text{O-CHR}^{18}\text{CHR}^{19})_h$, $(\text{CHR}^{18}\text{CHR}^{19}\text{-O})_h$, C=O, C=S, C=NR¹⁵, $\text{CH}(\text{OR}^{15})$, $\text{C}(\text{OR}^{15})(\text{OR}^{20})$, $\text{C}(=\text{O})\text{O}$, $\text{OC}(=\text{O})$, $\text{OC}(=\text{O})\text{O}$, $\text{C}(=\text{O})\text{N}(\text{R}^{15})$, $\text{N}(\text{R}^{15})\text{C}(=\text{O})$, $\text{CH}=\text{N-O}$, $\text{CH}=\text{N-NR}^{15}$, $\text{OC}(\text{O})\text{NR}^{15}$, $\text{NR}^{15}\text{C}(\text{O})\text{O}$, S=O, SO₂, SO₂NR¹⁵ and NR¹⁵SO₂, wherein
- are independently from each other 0, 1, 2, 3, 4, 5, or 6, and
- is 1, 2, 3, 4, 5, or 6,
- is selected from O, S, NR²¹, $\text{C}(\text{R}^{22})\text{-NO}_2$, $\text{C}(\text{R}^{22})\text{-CN}$ and $\text{C}(\text{CN})_2$, wherein
- is independently selected from the meanings given for R¹³, R¹⁴ and

- are independently selected from a group consisting of H, A, $(\text{CH}_2)_m\text{Ar}^3$ and $(\text{CH}_2)_m\text{Het}$, or in NR¹¹R¹²,
- form, together with the N-atom they are bound to, a 5-, 6- or 7- membered heterocycly which optionally contains 1 or 2 additional hetero atoms, selected from N, O and S,
- are independently selected from a group consisting of H, Hal, A, $(\text{CH}_2)_m\text{Ar}^4$ and $(\text{CH}_2)_m\text{Het}$,
- is selected from the group consisting of alkyl, alkenyl, cycloalkyl, alkenecycloalkyl, alkoxy, alkoxyalkyl and saturated heterocyclyl, preferably from the group consisting of alkyl, alkenyl, cycloalkyl, alkenecycloalkyl, alkoxy and alkoxyalkyl,
- are independently from one another aromatic hydrocarbon residues comprising 5 to 12 and preferably 5 to 10 carbon atoms which are optionally substituted by one or more substituents, selected from a group consisting of A, Hal, NO₂, CN, OR¹⁵, NR¹⁵R¹⁶, COOR¹⁵, CONR¹⁵R¹⁶, NR¹⁵COR¹⁶, NR¹⁵CONR¹⁵R¹⁶, NR¹⁶SO₂A, COR¹⁵, SO₂R¹⁵⁻¹⁶, S(O)₂A and OOCR¹⁵,
- is a saturated, unsaturated or aromatic heterocyclic residue which is optionally substituted by one or more substituents, selected from a group consisting of A, Hal, NO₂, CN, OR¹⁵, NR¹⁵R¹⁶, COOR¹⁵, CONR¹⁵R¹⁶, NR¹⁵COR¹⁶, NR¹⁵CONR¹⁵R¹⁶, NR¹⁶SO₂A, COR¹⁵, SO₂R¹⁵⁻¹⁶, S(O)₂A and OOCR¹⁵,

- 44 -

Especially preferred as compounds of formula II are compounds of formula II',



10 wherein E and G are as defined above, preferably E and G are both nitrogen atoms; more preferably one of E and G is a nitrogen atom or both E and G are carbon atoms. If E and/or G are carbon atoms, they can be unsubstituted or substituted by R⁹, i. e. E and/or G are either CH or CR⁹.

15 In compounds of formula II, the term alkyl preferably refers to an unbranched or branched alkyl residue, preferably an unbranched alkyl residue comprising 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10, preferably 1, 2, 3, 4, 5 or 6, more preferred 1, 2, 3 or 4 and especially 1 or 2 carbon atoms, or a branched alkyl residue comprising 3, 4, 5, 6, 7, 8, 9 or 10, preferably 3, 4, 5 or 6 more preferred 3 or 4 carbon atoms. The alkyl residues can be optionally substituted, especially by one or more halogen atoms, for example up to perhaloalkyl, by one or more hydroxy groups or by one or more amino groups, all of which can optionally be substituted by alkyl. If an alkyl residue is substituted by halogen, it usually comprises 1, 2, 3, 4 or 5 halogen atoms, depending on the number of carbon atoms of the alkyl residue. For example, a methyl group can comprise, 1, 2 or 3 halogen atoms, an ethyl group (an alkyl residue comprising 2 carbon atoms) can comprise 1, 2, 3, 4 or 5 halogen atoms. If an alkyl residue is substituted by hydroxy groups, it usually comprises one or two, preferably one hydroxy groups. If the hydroxy group is substituted by alkyl, the alkyl substituent comprises preferably 1 to 4 carbon atoms and is preferably unsubstituted or substituted by halogen and more preferred unsubstituted. If an alkyl residue is substituted by amino groups, it usually comprises one or two,

- 45 -

5 preferably one amino groups. If the amino group is substituted by alkyl, the alkyl substituent comprises preferably 1 to 4 carbon atoms and is preferably unsubstituted or substituted by halogen and more preferred unsubstituted. According to compounds of formula II, alkyl is preferably selected from the group consisting of methyl, ethyl, trifluoro methyl, pentafluoro ethyl, isopropyl, tert.-butyl, 2-amino ethyl, N-methyl-2-amino ethyl, N,N-dimethyl-2-amino ethyl, N-ethyl-2-amino ethyl, N,N-diethyl-2-amino ethyl, 2-hydroxy ethyl, 2-methoxy ethyl and 2-ethoxy ethyl, further preferred of the group consisting of 2-butyl, n-pentyl, neo-nentyl, isopentyl, hexyl and n-decyl, more preferred of methyl, ethyl, trifluoro methyl, isopropyl and tert.-butyl.

10 In compounds of formula II, alkenyl is preferably selected from the group consisting of allyl, 2- or 3-butenyl, isobutenyl, sec-butenyl, furthermore preferably 4-pentenyl, isopentenyl and 5-hexenyl.

15 In compounds of formula II, alkylene is preferably unbranched and is more preferably methylene or ethylene, furthermore preferably propylene or butylene.

20 In compounds of formula II, alkylenecycloalkyl preferably has 5 to 10 carbon atoms and is preferably methylenecyclopropyl, methylenecyclobutyl, furthermore preferably methylenecyclopentyl, methylenecyclohexyl or methylenecycloheptyl, furthermore alternatively ethylenecyclopropyl, ethylenecyclobutyl, ethylenecyclopentyl, ethylenecyclohexyl or ethylenecycloheptyl, propylenecyclopentyl, propylenecyclohexyl, butylenecyclopentyl or butylenecyclohexyl.

30 In compounds of formula II, the term "alkoxy" preferably comprises groups of formula O-alkyl, where alkyl is an alkyl group as defined above. More preferred, alkoxy is selected from group consisting of methoxy, ethoxy, n-propoxy, isopropoxy, 2-butoxy, tert.-butoxy and halogenated, especially

perhalogenated, derivatives thereof. Preferred perhalogenated derivatives are selected from the group consisting of O-CCl_3 , O-CF_3 , $\text{O-C}_2\text{Cl}_5$, $\text{O-C}_2\text{F}_5$, $\text{O-C}(\text{CCl}_3)_3$ and $\text{O-C}(\text{CF}_3)_3$.

5 In compounds of formula II, the term "alkoxyalkyl" preferably comprises branched and unbranched residues, more preferred unbranched residues, of formula $\text{C}_u\text{H}_{2u+1}\text{O}-(\text{CH}_2)_v$, wherein u and v are independently from each other 1 to 6. Especially preferred is u = 1 and v 1 to 4.

10 In compounds of formula II the term "alkoxyalkyl" includes alkoxyalkyl groups as defined above, wherein one or more of the hydrogen atoms are substituted by halogen, for example up to perhalo alkoxyalkyl.

15 In compounds of formula II, cycloalkyl preferably has 3 – 7 carbon atoms and is preferably cyclopropyl or cyclobutyl, furthermore preferably cyclopentyl or cyclohexyl, furthermore also cycloheptyl, particularly preferably cyclopentyl. The term "cycloalkyl", as used herein preferably also includes saturated heterocyclic groups, wherein one or two carbon atoms are substituted by hetero atoms, selected from the group consisting of O, NH, NA and S, wherein A is as defined as above/below.

20 In compounds of formula II, Ar^3 to Ar^8 are preferably selected independently from one another from phenyl, naphthyl and biphenyl which is optionally substituted by one or more substituents, selected from the group consisting of A, Hal, NO_2 , CN, OR^{15} , $\text{NR}^{15}\text{R}^{16}$, COOR^{15} , $\text{CONR}^{15}\text{R}^{16}$, $\text{NR}^{15}\text{COR}^{16}$, $\text{NR}^{15}\text{CONR}^{15-16}$, $\text{NR}^{15}\text{SO}_2\text{A}$, COR^{15} , $\text{SO}_2\text{R}^{15-16}$, $\text{S}(\text{O})_n\text{A}$ and OOCR^{15} .

30 In compounds of formula II, het is preferably an optionally substituted aromatic heterocyclic residue and even more preferred and optionally substituted saturated heterocyclic residue, wherein the substituents are preferably selected from A, CN and hal. Even more preferred, het is

selected from the group consisting of 1-piperidyl, 1-piperazyl, 1-(4-methyl)-piperazyl, 4-methylpiperazin-1-yl amine, 4-morpholinyl, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-pyrazolidinyl 1-(2-methyl)-pyrazolidinyl, 1-imidazolidinyl or 1-(3-methyl)-imidazolidinyl, thiophen-2-yl, thiophen-3-yl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, chinolinyl, isochinolinyl, 2-pyridazyl, 4-pyridazyl, 2-pyrimidyl, 4-pyrimidyl, 5-pyrimidyl, 2-pyrazinyl and 3-pyrazinyl. Especially the thiophenyl and the pyridyl residues can optionally be substituted by one or more cyano groups.

10 In compounds of formula II, saturated heterocyclyl is preferably a substituted or unsubstituted saturated heterocyclic residue, more preferred an unsubstituted saturated heterocyclic residue, preferably selected from the saturated groups given above in the definition of het.

15 In compounds of formula II, aromatic hydrocarbons containing 6 to 14 carbon atoms and ethylenical unsaturated or aromatic heterocyclic residues containing 3 to 10 carbon atoms and one or two heteroatoms, independently selected from N, O and S, are preferably selected from the definitions given herein for aryl, heteroaryl and/or het. Heteroaryl is more preferably furanyl, thiophenyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, tetrazolyl, thiazolyl, oxazolyl, isoxazolyl, oxadiazolyl, oxo-pyridyl, thiadiazolyl, isothiazolyl, pyridyl, pyridazyl, pyrazinyl, pyrimidyl, quinolinyl, isoquinolinyl, benzofuranyl, benzothiophenyl, indolyl, indazolyl and even more preferably pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl and/or imidazolyl. Aryl more preferably refers to an optionally substituted benzene ring or to an optionally substituted benzene ring system fused to one or more optionally substituted benzene rings to form, for example,

25 anthracene, phenanthrene, or naphthalene ring systems. Even more preferably, aryl is selected from the group consisting of phenyl, 2-naphthyl, 1-naphthyl, biphenyl.

30

- 48 -

5 In compounds of formula II, Ar¹ is preferably selected from the group consisting of phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl and imidazolyl, and especially from phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl and oxazolyl.

10 In compounds of formula II, Ar² is preferably selected from the group consisting of phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl and imidazolyl, even more preferably from phenyl, pyridinyl and pyrimidyl and especially preferred from phenyl and pyridinyl.

15 Preferably, the sum of h and l exceeds 0.

A preferred aspect of the instant invention relates to compounds of formula II, wherein n is 0 or 1 and especially 0.

20 Another preferred aspect of the instant invention relates to compounds of formula II, wherein n is 0 in the residues R⁸, R⁹ and/or R¹⁰ and especially in R¹⁰.

25 Another preferred aspect of the instant invention relates to compounds of formula II, wherein n is 0 in the residues R⁸ and/or R⁷.

Another preferred aspect of the instant invention relates to compounds of formula II, wherein X represents a bridging group, selected from (CR¹¹R¹²)_n or (CHR¹¹)_n-Q-(CHR¹²)_n.

30

- 49 -

The invention relates in particular to compounds of the formula II in which at least one of said radicals has one of the preferred meanings given above.

5 Some more preferred groups of compounds may be expressed by the following sub-formulae II.1) to II.20), which correspond to the formula II and in which radicals not denoted in greater detail are as defined in the formula II, but in which

10 II.1) Ar¹ is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl;

15 II.2) Ar¹ is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl, and

p is 1, 2 or 3;

20 II.3) Ar¹

is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,

25

30 p is 1, 2 or 3, and

- 50 -

- R⁸** is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_kR¹³;
- II.4) Ar¹** is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
- p** is 1, 2 or 3,
- R⁸** is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_kR¹³,
- II.5) Ar¹** is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,

- 51 -

- p** is 1, 2 or 3,
- R⁸** is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_kR¹³, wherein
- n** is 0 or 1;
- Ar¹** is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
- p** is 1, 2 or 3,
- R⁸** is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_kR¹³, wherein

- 52 -

	n	is 0 or 1, and
	u	is 0;
5	II.7) Ar ¹	is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
10	p	is 1, 2 or 3,
15	R ⁸	is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH ₂ Hal, CH(Hal) ₂ , perhaloalkyl comprising 1 to 4 carbon atoms, NO ₂ , (CH ₂) _n CN, (CH ₂) _n NR ¹ R ¹² , (CH ₂) _n O(CH ₂) _n NR ¹ R ¹² , (CH ₂) _n NR ¹ (CH ₂) _n R ¹² , (CH ₂) _n O(CH ₂) _n OR ¹ , (CH ₂) _n NR ¹ (CH ₂) _n OR ¹² , (CH ₂) _n COOR ¹³ , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹ R ¹² , (CH ₂) _n SO ₂ NR ¹ R ¹² and (CH ₂) _n S(O) _n R ¹³ , wherein
20	n	is 0 or 1,
	u	is 0, and
25	q	is 0 or 1, and
30	X	is selected from the group consisting of O, S, NR ¹ , CHOR ¹ , CH ₂ , CH ₂ CH ₂ , OCH ₂ , CH ₂ O, OCH ₂ CH ₂ , CH ₂ CH ₂ O, preferably O, S and CH ₂ and especially O and S;

- 53 -

	II.8) Ar ¹	is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
	p	is 1, 2 or 3,
10	R ⁸	is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH ₂ Hal, CH(Hal) ₂ , perhaloalkyl comprising 1 to 4 carbon atoms, NO ₂ , (CH ₂) _n CN, (CH ₂) _n NR ¹ R ¹² , (CH ₂) _n O(CH ₂) _n NR ¹ R ¹² , (CH ₂) _n NR ¹ (CH ₂) _n R ¹² , (CH ₂) _n O(CH ₂) _n OR ¹ , (CH ₂) _n NR ¹ (CH ₂) _n OR ¹² , (CH ₂) _n COOR ¹³ , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹ R ¹² , (CH ₂) _n SO ₂ NR ¹ R ¹² and (CH ₂) _n S(O) _n R ¹³ , wherein
15	15	
20	n	is 0 or 1,
	u	is 0, and
25	q	is 0 or 1, and
30	X	is selected from the group consisting of O, S, NR ¹ , CHOR ¹ , CH ₂ , CH ₂ CH ₂ , OCH ₂ , CH ₂ O, OCH ₂ CH ₂ , CH ₂ CH ₂ O, preferably O, S and CH ₂ and especially O and S,
30	Ar ²	is phenyl, pyridinyl or pyrimidyl, and especially is phenyl or pyridinyl;

II.9)	A ^{r1}	is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,	R ¹⁰	is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH ₂ Hal, CH(Hal) ₂ , perhaloalkyl comprising 1 to 4 carbon atoms, NO ₂ , (CH ₂) _n CN, (CH ₂) _n NR ¹¹ R ¹² , (CH ₂) _n O(CH ₂) _k NR ¹¹ R ¹² , (CH ₂) _n NR ¹¹ (CH ₂) _k NR ¹¹ R ¹² , (CH ₂) _n O(CH ₂) _k OR ¹¹ , (CH ₂) _n NR ¹¹ (CH ₂) _k OR ¹² , (CH ₂) _n COR ¹³ , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹¹ R ¹² , (CH ₂) _n SO ₂ NR ¹¹ R ¹² and (CH ₂) _n S(O) _u R ¹³ , preferably alkyl comprising 1 to 4 carbon atoms, (CH ₂) _n NR ¹¹ R ¹² , (CH ₂) _n O(CH ₂) _k NR ¹¹ R ¹² , (CH ₂) _n COR ¹³ , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹¹ R ¹² and especially (CH ₂) _n CONR ¹¹ R ¹² ,
5			5	
10	R ⁸	is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH ₂ Hal, CH(Hal) ₂ , perhaloalkyl comprising 1 to 4 carbon atoms, NO ₂ , (CH ₂) _n CN, (CH ₂) _n NR ¹¹ R ¹² , (CH ₂) _n O(CH ₂) _k NR ¹¹ R ¹² , (CH ₂) _n NR ¹¹ (CH ₂) _k OR ¹¹ , (CH ₂) _n O(CH ₂) _k OR ¹² , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹¹ R ¹² , (CH ₂) _n SO ₂ NR ¹¹ R ¹² and (CH ₂) _n S(O) _u R ¹³ , wherein	10	
15			15	II.10) Ar ¹
20	n	is 0 or 1,	20	is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl, pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
	u	is 0, and		is 1, 2 or 3,
	q	is 0 or 1, and		is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH ₂ Hal, CH(Hal) ₂ , perhaloalkyl comprising 1 to 4 carbon atoms, NO ₂ , (CH ₂) _n CN, (CH ₂) _n NR ¹¹ R ¹² , (CH ₂) _n O(CH ₂) _k NR ¹¹ R ¹² , (CH ₂) _n NR ¹¹ (CH ₂) _k OR ¹¹ , (CH ₂) _n O(CH ₂) _k OR ¹² , (CH ₂) _n COOR ¹³ , (CH ₂) _n CONR ¹¹ R ¹² , (CH ₂) _n SO ₂ NR ¹¹ R ¹² and (CH ₂) _n S(O) _u R ¹³ , wherein
25	X	is selected from the group consisting of O, S, NR ¹¹ , CHOR ¹¹ , CH ₂ , CH ₂ CH ₂ , OCH ₂ , CH ₂ O, OCH ₂ CH ₂ , CH ₂ CH ₂ O, preferably O, S and CH ₂ and especially O and S,	25	
30	Ar ²	is phenyl, pyridinyl or pyrimidyl, and especially is phenyl or pyridinyl; and	30	n is 0 or 1,

- 56 -

- u is O, and
- q is 0 or 1, and
- 5 X is selected from the group consisting of O, S, NR¹¹, CHOR¹¹, CH₂, CH₂CH₂, OCH₂, CH₂O, OCH₂CH₂, CH₂CH₂O, preferably O, S and CH₂ and especially O and S,
- 10 Ar² is phenyl, pyridinyl or pyrimidyl, and especially is phenyl or pyridinyl; and
- R¹⁰ is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_nNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_nKOR¹¹, (CH₂)_nNR¹¹(CH₂)_nNR¹¹R¹², (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², and (CH₂)_nCONR¹¹R¹², preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_nNR¹¹R¹², (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹² and especially (CH₂)_nCONR¹¹R¹², wherein
- 25 n is 0, 1 or 2, preferably 0 or 1;
- II.11) Ar¹ is phenyl, pyridinyl, pyrimidyl, chinolinyl, isochinolinyl, thiophenyl, thiadiazolyl, benzothiadiazolyl, oxazolyl, isoxazolyl, pyrazolyl or imidazolyl, preferably phenyl,
- 30

- 57 -

- pyridinyl, chinolinyl, isochinolinyl, thiophenyl, benzothiadiazolyl, oxazolyl, isoxazolyl or oxazolyl,
- p is 1, 2 or 3,
- 5 R⁸ is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_nNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_nKOR¹¹, (CH₂)_nNR¹¹(CH₂)_nOR¹², (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)R¹³, wherein
- 10 n is 0 or 1,
- u is 0, and
- q is 0 or 1, and
- 20 X is selected from the group consisting of O, S, NR¹¹, CHOR¹¹, CH₂, CH₂CH₂, OCH₂, CH₂O, OCH₂CH₂, CH₂CH₂O, preferably O, S and CH₂ and especially O and S,
- 25 Ar² is phenyl, pyridinyl or pyrimidyl, and especially is phenyl or pyridinyl; and
- R¹⁰ is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl
- 30

comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹² and (CH₂)_nS(O)_nR¹³, preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹² and especially (CH₂)_nCONR¹¹R¹², wherein

is 0, 1 or 2, preferably 0 or 1 and

is 0, 1 or 2, preferably 0 or 1;

is 1, 2 or 3,

is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_nR¹³, wherein

is 0 or 1,

is 0, and

is 0 or 1, and

is selected from the group consisting of O, S, NR¹¹, CHOR¹¹, CH₂, CH₂CH₂, OCH₂, CH₂O, OCH₂CH₂, CH₂CH₂O, preferably O, S and CH₂ and especially O and S,

is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl or pyridinyl; and

is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹², (CH₂)_nSO₂NR¹¹R¹² and (CH₂)_nS(O)_nR¹³, preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹² and especially (CH₂)_nCONR¹¹R¹², wherein

is 0, 1 or 2, preferably 0 or 1 and

is 0, 1 or 2, preferably 0 or 1;

is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹¹R¹², (CH₂)_nO(CH₂)_kNR¹¹R¹², (CH₂)_nNR¹¹(CH₂)_kNR¹¹R¹², (CH₂)_nO(CH₂)_kOR¹¹, (CH₂)_nNR¹¹(CH₂)_kOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹¹R¹² and especially (CH₂)_nCONR¹¹R¹², wherein

30

X

5

Al²

R¹⁰

10

15

20

n

r

25

Il.13) R⁸.

30

- 60 -

$(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$,
 $(CH_2)_nSO_2NR^{11}R^{12}$ and $(CH_2)_nS(O)_nR^{13}$, wherein

n Is 0 or 1,

u Is 0, and

q Is 0 or 1, and

10 X Is selected from the group consisting of O, S, NR^{11} ,
 $CHOR^{11}$, CH_2 , CH_2CH_2 , OCH_2 , CH_2O , OCH_2CH_2 ,
 CH_2CH_2O , preferably O, S and CH_2 and especially O
 and S,

15 A^2 Is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl
 or pyridinyl; and

R^{10} Is selected from the group consisting of H, alkyl
 comprising 1 to 4 carbon atoms, alkoxy comprising 1 to
 4 carbon atoms, Hal, CH_2Hal , $CH(Hal)_2$, perhaloalkyl
 comprising 1 to 4 carbon atoms, NO_2 , $(CH_2)_nCN$,
 $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nNR^{11}(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nOR^{11}$,
 $(CH_2)_nNR^{11}(CH_2)_nOR^{12}$, $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$,
 $(CH_2)_nCONR^{11}R^{12}$, $(CH_2)_nSO_2NR^{11}R^{12}$ and
 $(CH_2)_nCONR^{11}R^{12}$, preferably alkyl comprising 1 to 4
 carbon atoms, $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$ and
 especially $(CH_2)_nCONR^{11}R^{12}$, wherein

30 n Is 0, 1 or 2, preferably 0 or 1 and

- 61 -

r Is 0, 1 or 2, preferably 0 or 1;

5 II.14) R^8 Is selected from the group consisting of alkyl comprising
 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon
 atoms, Hal, CH_2Hal , $CH(Hal)_2$, perhaloalkyl comprising
 1 to 4 carbon atoms, NO_2 , $(CH_2)_nCN$, $(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nNR^{11}(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nO(CH_2)_nOR^{11}$, $(CH_2)_nNR^{11}(CH_2)_nOR^{12}$,
 $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$,
 $(CH_2)_nSO_2NR^{11}R^{12}$ and $(CH_2)_nS(O)_nR^{13}$, wherein

u Is 0, and

15 q Is 0 or 1, and

X Is selected from the group consisting of O, S, NR^{11} ,
 $CHOR^{11}$, CH_2 , CH_2CH_2 , OCH_2 , CH_2O , OCH_2CH_2 ,
 CH_2CH_2O , preferably O, S and CH_2 and especially O
 and S,

A^2 Is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl
 or pyridinyl; and

25 R^{10} Is selected from the group consisting of H, alkyl
 comprising 1 to 4 carbon atoms, alkoxy comprising 1 to
 4 carbon atoms, Hal, CH_2Hal , $CH(Hal)_2$, perhaloalkyl
 comprising 1 to 4 carbon atoms, NO_2 , $(CH_2)_nCN$,
 $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nNR^{11}(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nOR^{11}$,
 $(CH_2)_nNR^{11}(CH_2)_nOR^{12}$, $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$,
 $(CH_2)_nCONR^{11}R^{12}$ and $(CH_2)_nSO_2NR^{11}R^{12}$, preferably alkyl
 comprising 1 to 4 carbon atoms, $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$,
 $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$ and
 especially $(CH_2)_nCONR^{11}R^{12}$, wherein

30

$(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{SO}_2\text{NR}^{11}\text{R}^{12}$ and $(\text{CH}_2)_n\text{S(O)}_n\text{R}^{13}$, preferably alkyl comprising 1 to 4 carbon atoms, $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{COR}^{13}$, $(\text{CH}_2)_n\text{COOR}^{13}$, $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$ and especially $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, wherein

5

is 0, 1 or 2, preferably 0 or 1 and

n

is 0, 1 or 2, preferably 0 or 1;

r

is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH_2Hal , $\text{CH}(\text{Hal})_2$, perhaloalkyl comprising 1 to 4 carbon atoms, NO_2 , $(\text{CH}_2)_n\text{CN}$, $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{OR}^{11}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{OR}^{12}$, $(\text{CH}_2)_n\text{COR}^{13}$, $(\text{CH}_2)_n\text{COOR}^{13}$, $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{SO}_2\text{NR}^{11}\text{R}^{12}$ and $(\text{CH}_2)_n\text{S(O)}_n\text{R}^{13}$, wherein

15

is 0 or 1, and

q

is selected from the group consisting of O, S, NR^{11} , CHOR^{11} , CH_2 , CH_2CH_2 , OCH_2 , CH_2O , OCH_2CH_2 , $\text{CH}_2\text{CH}_2\text{O}$, preferably O, S and CH_2 and especially O and S,

20

X

is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl or pyridinyl; and

25

 Ar^2

is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH_2Hal , $\text{CH}(\text{Hal})_2$, perhaloalkyl

30

 R^{10}

comprising 1 to 4 carbon atoms, NO_2 , $(\text{CH}_2)_n\text{CN}$, $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{OR}^{11}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{OR}^{12}$, $(\text{CH}_2)_n\text{COR}^{13}$, $(\text{CH}_2)_n\text{COOR}^{13}$, $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{SO}_2\text{NR}^{11}\text{R}^{12}$ and $(\text{CH}_2)_n\text{S(O)}_n\text{R}^{13}$, preferably alkyl comprising 1 to 4 carbon atoms, $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{COR}^{13}$, $(\text{CH}_2)_n\text{COOR}^{13}$, $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$ and especially $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, wherein

5

is 0, 1 or 2, preferably 0 or 1 and

n

is 0, 1 or 2, preferably 0 or 1;

r

is 0 or 1, and

15

II.16) q

is selected from the group consisting of O, S, NR^{11} , CHOR^{11} , CH_2 , CH_2CH_2 , OCH_2 , CH_2O , OCH_2CH_2 , $\text{CH}_2\text{CH}_2\text{O}$, preferably O, S and CH_2 and especially O and S,

20

is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl or pyridinyl; and

 Ar^2

is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH_2Hal , $\text{CH}(\text{Hal})_2$, perhaloalkyl comprising 1 to 4 carbon atoms, NO_2 , $(\text{CH}_2)_n\text{CN}$, $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{NR}^{11}\text{R}^{12}$, $(\text{CH}_2)_n\text{O}(\text{CH}_2)_k\text{OR}^{11}$, $(\text{CH}_2)_n\text{NR}^{11}(\text{CH}_2)_k\text{OR}^{12}$, $(\text{CH}_2)_n\text{COR}^{13}$, $(\text{CH}_2)_n\text{COOR}^{13}$, $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$ and especially $(\text{CH}_2)_n\text{CONR}^{11}\text{R}^{12}$, wherein

25

 R^{10}

30

- 64 -

- (CH₂)_nCONR¹R¹², (CH₂)_nSO₂NR¹R¹² and (CH₂)_nS(O)_nR¹³, preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹² and especially (CH₂)_nCONR¹R¹², wherein

n is 0, 1 or 2, preferably 0 or 1 and

r is 0, 1 or 2, preferably 0 or 1;

- 10 II.17) X is selected from the group consisting of O, S, NR¹, CHOR¹, CH₂, CH₂CH₂, OCH₂, CH₂O, OCH₂CH₂, CH₂CH₂O, preferably O, S and CH₂ and especially O and S,

- 15 A² is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl or pyridinyl; and

- 20 R¹⁰ is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nNR¹(CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nOR¹¹, (CH₂)_nNR¹(CH₂)_nOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹², (CH₂)_nSO₂NR¹R¹² and (CH₂)_nS(O)_nR¹³, preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹² and especially (CH₂)_nCONR¹R¹², wherein

- 65 -

- n is 0, 1 or 2, preferably 0 or 1 and
r is 0, 1 or 2, preferably 0 or 1;

- 5 II.18) A² is phenyl, pyridinyl or pyrimidinyl, and especially is phenyl or pyridinyl; and

- 10 R¹⁰ is selected from the group consisting of alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nNR¹(CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nOR¹, (CH₂)_nNR¹(CH₂)_nOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹², (CH₂)_nSO₂NR¹R¹² and (CH₂)_nS(O)_nR¹³, preferably alkyl comprising 1 to 4 carbon atoms, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nNR¹(CH₂)_nOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹² and especially (CH₂)_nCONR¹R¹²,

- 20 n is 0, 1 or 2, preferably 0 or 1 and

r is 0, 1 or 2, preferably 0 or 1;

- 25 II.19) R¹⁰ is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH₂Hal, CH(Hal)₂, perhaloalkyl comprising 1 to 4 carbon atoms, NO₂, (CH₂)_nCN, (CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nNR¹R¹², (CH₂)_nNR¹(CH₂)_nNR¹R¹², (CH₂)_nO(CH₂)_nOR¹¹, (CH₂)_nNR¹(CH₂)_nOR¹², (CH₂)_nCOR¹³, (CH₂)_nCOOR¹³, (CH₂)_nCONR¹R¹², (CH₂)_nSO₂NR¹R¹² and

$(CH_2)_nS(O)_pR^{13}$, preferably alkyl comprising 1 to 4 carbon atoms, $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kNR^{11}R^{12}$, $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$ and especially $(CH_2)_nCONR^{11}R^{12}$,

5

n is 0, 1 or 2, preferably 0 or 1 and

r is 0, 1 or 2, preferably 0 or 1;

10 II.20) R^{10}

is selected from the group consisting of H, alkyl comprising 1 to 4 carbon atoms, alkoxy comprising 1 to 4 carbon atoms, Hal, CH_2Hal , $CH(Hal)_2$, perhaloalkyl

comprising 1 to 4 carbon atoms, NO_2 , $(CH_2)_nCN$,

$(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kNR^{11}R^{12}$,

$(CH_2)_nNR^{11}(CH_2)_kNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kOR^{11}$,

$(CH_2)_nNR^{11}(CH_2)_kOR^{12}$, $(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$,

$(CH_2)_nCONR^{11}R^{12}$, $(CH_2)_nSO_2NR^{11}R^{12}$ and

$(CH_2)_nS(O)_pR^{13}$, preferably alkyl comprising 1 to 4

carbon atoms, $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kNR^{11}R^{12}$,

$(CH_2)_nCOR^{13}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$ and

especially $(CH_2)_nCONR^{11}R^{12}$, and

20

r is 0, 1 or 2, preferably 0 or 1.

25 One preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein p is 1, 2 or 3 and R^8 is independently selected from the group consisting of methyl, ethyl, isopropyl, tert-butyl, F, Cl, Br, CF_3 , $C(CF_3)_3$, SO_2CF_3 , methoxy, ethoxy, tert-butoxy, perfluoro tert-butoxy ($OC(CF_3)_3$), methyl sulfanyl (SCH_3), ethyl sulfanyl (SCH_2CH_3), acetyl ($COCH_3$),

30

propionyl ($COCH_2CH_3$), butyryl ($COCH_2CH_2CH_3$). If p is 2 or 3, all substituents can be the same or different.

Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein X is selected from the group consisting of S, NR^{21} , CH_2 , CH_2CH_2 , OCH_2 and CH_2O .

5

Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein X is selected from the group consisting of S, CH_2 .

10

Another even more preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein X is O.

15

Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Y is selected from the group consisting of $C(R^{22})NO_2$, $C(R^{22})CN$ and $C(CN)_2$.

20

Another more preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Y is selected from the group consisting of O, S and NR^{21} .

25

Another even more preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Y is selected from the group consisting of O and S.

30 Another even more preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Y is O.

- 68 -

Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein R^6 and R^7 both are hydrogen.

- 5 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein R^8 or R^7 is a residue other than hydrogen. In this embodiment, the residue other than hydrogen is preferably selected from the meanings given for R^8 , R^9 and R^{10} , more preferably from A, and especially preferred from substituted or preferably unsubstituted alkyl, substituted or preferably unsubstituted alkenyl, substituted or preferably unsubstituted cycloalkyl and substituted or preferably unsubstituted alkenecycloalkyl, even more preferred substituted or unsubstituted alkyl with 1 to 6 carbon atoms, for example methyl, ethyl, n-propyl, isopropyl, n-butyl, 2-butyl, tert.-butyl, optionally substituted by one or more hydroxy groups, preferably one or two hydroxy groups and/or one or more halogen atoms, up to perhalo. Examples for preferred substituted alkyl groups are CH_2Hal , especially CH_2F , CH_2Cl and CH_2Br , CH_2I , especially CF_3 , CCl_3 and CBR_3 , and $(CH_2)_nOH$, wherein Z is 1 to 6, especially CH_2OH and CH_2CH_2OH .
- 10
- 15
- 20

- Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein R^6 or R^7 is a residue other than hydrogen. In this embodiment, the residue other than hydrogen is preferably selected from $(CH_2)_nCOOR^{13}$, $(CH_2)_nCONR^{11}R^{12}$, $(CH_2)_nNR^{11}COR^{13}$, $(CH_2)_nNR^{11}CONR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$ and $(CH_2)_nNR^{11}(CH_2)_nNR^{11}R^{12}$, more preferred $(CH_2)_nCOOR^{13}$, even more preferred $(CH_2)_nCOOA$ and $(CH_2)_nCOOH$ and especially preferred
- 25
- 30

- 69 -

$(CH_2)_nCOOA$, wherein A is C_1-C_n -alkyl and $(CH_2)_nCOOH$. In this embodiment, n is as defined above/below and especially is 0.

- 5 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein R^8 is hydrogen and R^7 is methyl, or R^8 is methyl and R^7 is hydrogen.

- 10 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein R^8 and R^7 are residues other than hydrogen. In this embodiment, R^8 and R^7 are preferably selected, independently from one another, from the meanings given for R^8 , R^9 and R^{10} , more preferably from the meanings given for A, and especially preferred from substituted or preferably unsubstituted alkyl, substituted or preferably unsubstituted alkenyl, substituted or preferably unsubstituted cycloalkyl and substituted or preferably unsubstituted alkenecycloalkyl, even more preferred substituted or unsubstituted alkyl with 1 to 6 carbon atoms, for example methyl, ethyl, n-propyl, isopropyl, n-butyl, 2-butyl, tert.-butyl, optionally substituted by one or more hydroxy groups, preferably one or two hydroxy groups and/or one or more halogen atoms, up to perhalo. Examples for preferred substituted alkyl groups are CH_2Hal , especially CH_2F , CH_2Cl and CH_2Br , CH_2I , especially CF_3 , CCl_3 and CBR_3 , and $(CH_2)_nOH$, wherein Z is 1 to 6, especially CH_2OH and CH_2CH_2OH . In this embodiment, R^8 and R^7 even more preferred form, together with the carbon atom they are bound to (i. e. the carbon atom of the methylene moiety of the methylene urea moiety), a carbocyclic residue comprising 3 to 6 carbon atoms or a heterocyclic residue comprising one or two heteroatoms, selected from the group consisting of O, N and S, and 2 to 5 carbon atoms, wherein the carbocyclic residue respectively the heterocyclic residue can be substituted by one or more substituents, preferably one or two substituents, selected, independently from one another, from the
- 15
- 20
- 25
- 30

meanings given for R^8 , R^9 and R^{10} . If R^6 and R^7 form a cyclic residue together with the carbon atom they are bound to, carbocyclic residues are preferred. Even more preferred are carbocyclic residues comprising 3, 4 or 5 carbon atoms, especially 3 carbon atoms which can be substituted once or twice as given above and preferably are unsubstituted. In this respect, one preferred embodiment of the instant invention relates to compounds, wherein R^6 and R^7 form, together with the carbon atom they are bound to, a cyclopropane moiety.

10 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein one of the residues R^6 or R^7 or both residues R^6 and R^7 are other than hydrogen and are preferably as defined in the preferred embodiments relating to R^6 and R^7 given above.

15 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar^2 is pyridinyl.

20 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein r is either 0 or 1. If r is 1, R^{10} is preferably $(CH_2)_nCONR^{11}R^{12}$ and especially $(CH_2)_nCONR^{11}R^{12}$, wherein n in 0. In this embodiment, R^{11} is preferably selected from the group consisting of H and A and more preferred from H and alkyl, and R^{12} is preferably selected from the group consisting of H and A and more preferred from H and alkyl.

25 Especially preferred as residue R^{10} are carbamoyl, more preferred alkyl carbamoyl or dialkyl carbamoyl, even more preferred methyl carbamoyl or dimethyl carbamoyl, ethyl carbamoyl or diethyl carbamoyl and especially preferred methyl carbamoyl (-CONHCH₃). This embodiment is especially preferred when Ar^2 is pyridinyl. When Ar^2 is pyridinyl, R^{10} is preferably

30

bonded in a vicinal position to the nitrogen atom of the pyridinyl residue, i.e. in 2- and/or 6-position of the pyridinyl residue.

5 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar^1 comprises two or more substituents R^8 , wherein one or more, preferably one substituent R^8 is selected from the group consisting of $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kNR^{11}R^{12}$, $(CH_2)_nNR^{11}(CH_2)_kOR^{12}$, $(CH_2)_nNR^{11}(CH_2)_kNR^{12P^{12}}$, $(CH_2)_nCOOR^{13}$ and $(CH_2)_nS(O)_uR^{13}$ wherein R^{11} , R^{12} and R^{13} are defined as above and n is as defined above, preferably n is 0, 1 or 2 and especially is 0, k is 1 to 4 and preferably 1 or 2, and u is preferably 2. In this embodiment R^{11} , R^{12} and R^{13} are more preferably selected independently from each other from the

10 group consisting of H, methyl and ethyl. In this embodiment, one or two substituents R^8 and preferably one substituent R^8 is especially preferably selected from the group consisting of NH_2 , $N(CH_3)_2$, $N(C_2H_5)_2$, $NHCH_2CH_2NH_2$, $N(CH_3)CH_2CH_2NH_2$, $N(CH_3)CH_2CH_2N(CH_3)_2$, $N(CH_3)CH_2CH_2N(CH_3)_2$, $N(CH_3)CH_2CH_2OCH_3$, $OCH_2CH_2N(CH_3)_2$, SCH_3 , SC_2H_5 , SO_2CH_3 , $COOCH_3$ and $COOH$. Accordingly, in this embodiment Ar^1 especially preferably comprises at least one substituent R^8 other than $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_kNR^{11}R^{12}$, $(CH_2)_nNR^{11}(CH_2)_kOR^{12}$, $(CH_2)_nNR^{11}(CH_2)_kNR^{12P^{12}}$, $(CH_2)_nCOOR^{13}$ and $(CH_2)_nS(O)_uR^{13}$ as defined in this paragraph and especially other than NH_2 , $N(CH_3)_2$, $N(C_2H_5)_2$, $NHCH_2CH_2NH_2$, $N(CH_3)CH_2CH_2NH_2$, $N(CH_3)CH_2CH_2N(CH_3)_2$, SCH_3 , SC_2H_5 , SO_2CH_3 , $COOCH_3$ and $COOH$.

20 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein q is 0, i.e. the phenyl group bound to the methylene group of the methylene urea moiety is unsubstituted.

30 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein q is 0, i.e. the phenyl group bound to the methylene group of the methylene urea moiety is unsubstituted.

- 72 -


Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein q is 1, i.e. the phenyl group bound to the methylene group of the methylene urea moiety is substituted by one substituent, preferably a substituent as defined above and more preferably a substituent selected from alkyl and hal, and especially selected from CH_3 , CH_2CH_3 and hal.

Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of formulae II.1) to II.20), wherein $(\text{R}^8)_p\text{-Ar}^1$ is selected from the group consisting of 3-acetyl-phenyl, 4-acetyl-phenyl, 2-bromo-phenyl, 3-bromo-phenyl, 4-bromo-phenyl, 4-bromo-2-chloro-phenyl, 4-bromo-3-methyl-phenyl, 4-bromo-3-trifluoromethyl-phenyl, 2-chloro-phenyl, 2-chloro-4-trifluoromethyl-phenyl, 2-chloro-5-trifluoromethyl-phenyl, 3-chloro-phenyl, 3-chloro-4-methyl-phenyl, 3-chloro-4-methoxy-phenyl, 3-chloro-4-methoxy-phenyl, 4-chloro-phenyl, 4-chloro-2-trifluoromethyl-phenyl, 4-chloro-3-trifluoromethyl-phenyl, 4-chloro-2-methyl-phenyl, 5-chloro-2-methyl-phenyl, 5-chloro-2-methoxy-phenyl, 2,3-dichloro-phenyl, 2,4-dichloro-phenyl, 2,5-dichloro-phenyl, 3,4-dichloro-phenyl, 3,5-dichloro-phenyl, 2,4,5-trichloro-phenyl, 4-fluoro-phenyl, 4-fluoro-3-trifluoromethyl-phenyl, 4-ethoxy-phenyl, 2-methoxy-phenyl, 2-methoxy-5-trifluoromethyl-phenyl, 4-methoxy-phenyl, 2,5-dimethoxy-phenyl, 2-trifluoromethyl-phenyl, 3-trifluoromethyl-phenyl, 3-trifluoromethoxy-phenyl, 4-trifluoromethyl-phenyl, 4-trifluoromethoxy-phenyl, 3,5-bis-trifluoromethyl-phenyl, 3-methoxy-phenyl, 3-methylsulfanyl-phenyl, 4-methylsulfanyl-phenyl, o-tolyl (2-methyl-phenyl), m-tolyl (3-methyl-phenyl), p-tolyl (4-methyl-phenyl), 2,3-dimethyl-phenyl, 2,5-dimethyl-phenyl, 3,4-dimethyl-phenyl, 3,5-dimethyl-phenyl, 2-ethyl-phenyl, 3-ethyl-phenyl, 4-ethyl-phenyl, 4-isopropyl-phenyl, 4-tert-butyl-phenyl and 5-tert-butyl-isoxazol-3-yl.


- 73 -

Another preferred embodiment of the instant invention relates to compounds of formula II and the subformulae related thereto and preferably one or more of formulae II.1) to II.20), wherein the residues $(\text{R}^8)_p\text{-Ar}^1$ are selected from the group consisting of compounds of the following formulae:


a)









- 74 -




- 75 -







- 78 -




5



and/or




10




15

and/or



20


and/or





25

30



- 79 -




5



10



15



and/or residues of the structures given above that comprise one or two, preferably one additional substituent, independently selected from the meanings given for R^8 .

20


Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein $(R^8)_{p-Ar^1}$ is as defined above, but comprises one or more additional residues, preferably one additional residue. The additional residues are preferably selected from the meanings given for R^8 and more preferably selected from the group consisting of $(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nO(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nNR^{11}(CH_2)_nOR^{12}$, $(CH_2)_nNR^{11}(CH_2)_nNR^{11}R^{12}$, $(CH_2)_nCOOR^{13}$, $(CH_2)_nS(O)_nR^{13}$ and $(CH_2)_nS(O)_nR^{13}$ wherein R^{11} , R^{12} and R^{13} are defined as above and n is as defined above, preferably n is 0,

25

30

1 or 2 and especially is 0, k is 1 to 4 and preferably 1 or 2, and u is preferably 2. In this embodiment R^{11} , R^{12} and R^{13} are more preferably selected independently from each other from the group consisting of H, methyl and ethyl. Even more preferred, the additional residue(s) s is/are selected from the group consisting of NH_2 , $N(CH_3)_2$, $N(C_2H_5)_2$, $NHC_2H_4CH_2NH_2$, $N(CH_3)CH_2CH_2NH_2$, $N(CH_3)CH_2CH_2N(CH_3)_2$, $N(CH_3)CH_2C_2H_4N(CH_3)_2$, $N(CH_3)CH_2C_2H_4OCH_3$, $OCH_2CH_2N(CH_3)_2$, SC_2H_5 , SO_2CH_3 , SO_2CF_3 , OSO_2CH_3 , OSO_2CF_3 , SO_2NH_2 , $SO_2NHCH(CH_3)_2$, $SO_2N(CH_3)_2$, $SO_2N(CH_2CH_3)_2$, 4-Morpholino-sulfonyl, $COOCH_3$ and $COOH$.

Another preferred embodiment of the instant invention relates to compounds of formula II and the subformulae related thereto and preferably one or more of formulae II.1) to II.20), wherein the residues $\text{Ar}^2\text{-(R}^{10})_n$ are selected from the group consisting of compounds of the following formulae:





and/or residues of the structures given above that comprise one or two, preferably one additional substituent, independently selected from the meanings given for R¹⁰.



5 Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein X is bonded in the para- (p-) or meta- (m-) position to the phenyl residue that is bonded directly to the methylene group of the methylene urea moiety.

Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein A^2 is a pyridinyl residue and wherein said pyridinyl residue is bonded to X in the 3- or 4-position, preferably the 4-position, relative to the nitrogen atom of the pyridinyl residue.


Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar¹ comprises one or more substituents R⁸ and wherein one or two, preferably one substituent R⁸ is selected from the group consisting of NH₂, N(CH₃)₂, N(C₂H₅)₂, NHCH₂CH₂NH₂, N(CH₃)CH₂CH₂NH₂, N(CH₃)CH₂CH₂N(CH₃)₂, N(CH₃)CH₂CH₂N(CH₃)₂, N(CH₃)CH₂CH₂OCH₃, OCH₂CH₂N(CH₃)₂, SCH₃, SC₂H₅, SO₂CH₃, SO₂CF₃, OSO₂CH₃, OSO₂CF₃, SO₂NH₂, SO₂NHCH(CH₃)₂, SO₂N(CH₃)₂, SO₂N(CH₂CH₃)₂, 4-Morpholinosulfonyl, COOCH₃ and COOH, more preferably NH₂, N(CH₃)₂, NHCH₃, N(C₂H₅)₂, HNCH₂CH₂NH₂, OCH₂CH₂NH₂, HOCH₂CH₂NH₂, OCH₂CH₂NHCH₃, N(CH₃)CH₂CH₂NH₂, HN(CH₃)CH₂CH₂NH₂, N(CH₃)CH₂CH₂N(CH₃)₂, N(CH₃)CH₂CH₂N(CH₃)₂, N(CH₃)CH₂CH₂OCH₃, OCH₂CH₂N(CH₃)₂, OCH₂CH₂N(CH₂CH₃)₂, SCH₃, SC₂H₅, and/or compounds of the formulae



5




10




and/or Ar² comprises one or more substituents R¹⁰ and wherein one or two, preferably one substituent R¹⁰ is independently selected from the meanings given for R⁸ in this paragraph.

15


Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar¹ comprises one or more substituents R⁸ and wherein one or two, preferably one substituent R⁸ is selected from the group consisting of



25



30



Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar¹ comprises one or more substituents R⁸ and wherein one or two, preferably one substituent R⁸ is selected from the group consisting of SO₂CH₃, SO₂CF₃, OSO₂CH₃, OSO₂CF₃, SO₂NH₂, SO₂NHCH(CH₃)₂, SO₂N(CH₃)₂, SO₂N(CH₂CH₃)₂ and 4-Morpholine-4-sulfonyl.

5

Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar² comprises one or more substituents R¹⁰ and wherein one or two, preferably one substituent R¹⁰ is selected from unsubstituted or substituted carbamoyl moieties. Substituted carbamoyl moieties are

10

preferably selected from CONHR²³ or CONR²³R²⁴, preferably CONHR²³, wherein R²³ and R²⁴ are independently selected from the definitions given for R⁸, more preferably selected from alkyl, preferably methyl, ethyl, propyl and butyl, (CH₂)_nNR¹¹R¹² and (CH₂)_nOR¹², wherein R¹¹, R¹² and n are as defined above. In this embodiment, n is preferably not 0 and more

15

preferred 1 to 3 and especially 1 or 2. Preferred examples for R²³ are

20

selected from the group consisting of methyl, ethyl, CH₂CH₂NH₂,

CH₂CH₂N(CH₃)₂, CH₂CH₂N(CH₂CH₃)₂, CH₂CH₂OH, CH₂CH₂OCH₃ and CH₂CH₂OCH₂CH₃.

25

Another preferred embodiment of the instant invention relates to

compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar² comprises one or more substituents R¹⁰ and wherein

one or two, preferably one substituent R¹⁰ is selected from substituted

carbamoyl moieties. Substituted carbamoyl moieties are preferably

selected from CONHR²³, wherein R²³ is preferably unsubstituted C₁-C₄-


30

alkyl and especially methyl.

- 84 -

- Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar^2 comprises one or more substituents R^{10} and wherein one or two, preferably one substituent R^{10} is selected from substituted carbamoyl moieties. Substituted carbamoyl moieties are preferably selected from $CONHR^{23}$, wherein R^{23} is selected from $(CH_2)_nR^{11}R^{12}$ and $(CH_2)_nOR^{12}$, wherein R^{11} , R^{12} and n are as defined above. In this embodiment, n is preferably not 0 and more preferred 1 to 3 and especially 1 or 2. Preferred examples for R^{23} are selected from the group consisting of $CH_2CH_2NH_2$, $CH_2CH_2N(CH_3)_2$, $CH_2CH_2N(CH_2CH_3)_2$, CH_2CH_2OH , $CH_2CH_2OCH_3$ and $CH_2CH_2OCH_2CH_3$.

Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein Ar^2 (R^{10}) is selected from the formulae





wherein R^{10} , R^{23} and R^{24} are as defined above and below.

- 85 -


Another especially preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20), wherein one or more features of the above and below mentioned embodiments are combined in one compound.

- Subject of the present invention are therefore preferably compounds of formula II according to one or both of the formulae IIa and IIb,




wherein Ar^1 , R^6 , p , Y , R^6 , R^7 , X , R^9 , q , Ar^2 , R^{10} and r are as defined above and below, and preferably as defined in sub formulae II.1) to II.20) and/or the embodiments related thereto.

Subject of the present invention are therefore especially preferred compounds of formula II according to one or both of the formulae IIc and IIId,




- 86 -




5

wherein Ar¹, R⁸, p, Y, R⁶, R⁷, X, R⁹ and q are as defined above and below, R¹⁰ is H or as defined above/below, and preferably as defined in sub formulae II.1) to II.20) and/or the embodiments related thereto;


10 and/or compounds of formula II according to one or more of the formulae IIe to IIx,



15




20




25

30


- 87 -




5




10




15



20








25








30

- 88 -




30

- 89 -



30



Iix

wherein R^6 , R^7 , R^8 , R^9 , p , Y , X , R^8 and q are as defined above and below, R^{10} is H or as defined above/below, and preferably as defined in sub formulae II.1) to II.20) and/or the embodiments related thereto.

One preferred aspect of the invention relates to compounds of formula II and especially to compounds of one or more of formulae IIa to Iix, wherein both R^6 and R^7 are hydrogen.

Another preferred aspect of the invention relates to compounds of formula II and especially to compounds of one or more of formulae IIa to Iix, wherein R^8 and/or R^7 are residues other than hydrogen.

Another preferred embodiment of the instant invention relates to compounds of formula II and preferably one or more of sub formulae II.1) to II.20) and IIa to Iix, wherein R^{10} is a substituted carbamoyl moiety CONHR^{23} or $\text{CONR}^{23}\text{R}^{24}$, preferably CONHR^{23} , wherein R^{23} and R^{24} are independently selected from the definitions given for R^8 , more preferably selected from $(\text{CH}_2)_n\text{NR}^{11}\text{R}^{12}$ and $(\text{CH}_2)_n\text{OR}^{12}$, wherein R^{11} , R^{12} and n are as defined above. In this embodiment, n is preferably not 0 and more preferred 1 to 3 and especially 1 or 2. Preferred examples for R^{23} are selected from the group consisting of $\text{CH}_2\text{CH}_2\text{NH}_2$, $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_3)_2$, $\text{CH}_2\text{CH}_2\text{N}(\text{CH}_2\text{CH}_3)_2$, $\text{CH}_2\text{CH}_2\text{OH}$, $\text{CH}_2\text{CH}_2\text{OCH}_3$ and $\text{CH}_2\text{CH}_2\text{OCH}_2\text{CH}_3$.

It is understood that when a residue, for example R^8 , R^9 , R^{10} or R^{14} or R^{23} , is comprised twice or more times in one or more of the formulae I, II and

















the sub formulae corresponding thereto, it is in each case independently from one another selected from the meanings given for the respective residue. For example, R^{11} and R^{12} are defined to be independently selected from a group consisting of H, A, $(\text{CH}_2)_m\text{Ar}^3$ and $(\text{CH}_2)_m\text{Het}$. Then $(\text{CH}_2)_m\text{NR}^{11}(\text{CH}_2)_m\text{NR}^{12}\text{R}^{12}$ can be $(\text{CH}_2)_m\text{NA}(\text{CH}_2)_m\text{NA}_2$ (if $R^{11} = A$, $R^{12} = A$ and $R^{12} = H$) as well as $(\text{CH}_2)_m\text{NA}(\text{CH}_2)_m\text{NHA}$ (if $R^{11} = A$, $R^{12} = H$ and $R^{12} = A$ or $(\text{CH}_2)_m\text{NA}(\text{CH}_2)_m\text{NH}(\text{CH}_2)_m\text{Het}$ (if $R^{11} = A$, $R^{12} = H$ and $R^{12} = (\text{CH}_2)_m\text{Het}$). Accordingly, if a compound of formula II comprises one residue R^8 , R^9 and R^{10} , then for example R^8 , R^9 and R^{10} can all be $(\text{CH}_2)_m\text{COOR}^{13}$, wherein all residues R^{13} are the same (for example CH_2Hal , wherein Hal is Cl; then all residues R^8 , R^9 and R^{10} are the same) or different (for example CH_2Hal , wherein in R^8 Hal is Cl; in R^9 Hal is F; and in R^{10} Hal is Br; then all residues R^8 , R^9 and R^{10} are different); or for example R^8 is $(\text{CH}_2)_m\text{COOR}^{13}$, R^9 is NO_2 and R^{10} is $(\text{CH}_2)_m\text{SR}^{11}$, wherein R^{11} and R^{13} can be the same (for example both can be H or both can be A which is methyl) of different (for example R^{11} can be H and R^{13} can be A which is methyl).

If not stated otherwise, reference to compounds of formula I and formula II also includes the sub formulae related thereto, especially sub formulae II.1) to II.20) and IIa to Iix.















Subject of the instant invention are especially those compounds of formula I and/or formula II, in which at least one of the residues mentioned in said formulae has one of the preferred or especially preferred meanings given above and below.

The present invention further relates to compounds (1) to (224) of formula $A\text{-NH-CO-NH-CH}_2\text{-B}$, wherein A and B are as given in the table below:


- 92 -

	A	B	Rt
(1)			2.12
(2)			4.67 ^a
(3)			4.10 ^a
(4)			3.87
(5)			2.43
(6)			2.46
(7)			1.77
(8)			1.80

- 93 -


(9)			1.66
(10)			2.43
(11)			1.93
(12)			1.95
(13)			2.65
(14)			2.65
(15)			1.94

- 94 -




5

1.95




10

2.70




15

2.68




20

2.02











25

2.72




30

2.75




- 95 -




5

2.30




10

1.59




15

2.30




20

2.30









25

2.71

















30













2.40




- 96 -

(29)			2.43	
5	(30)			1.45
10	(31)			2.23
15	(32)			2.21
20	(33)			1.56
25	(34)			1.55
30	(35)			2.31


- 97 -

(36)			2.32	
5	(37)			2.49
10	(38)			2.49
15	(39)			2.46
20	(40)			2.43
25	(41)			2.66
30				


- 98 -




2.65




1.88




2.55




2.55




1.57




2.29




2.29




2.65




1.88




2.55




2.55



1.57




2.29




2.29


- 99 -




1.82




1.81




2.48




2.47



1.47

















2.25















2.25


- 100 -

(56)			1.74
5 (57)			1.71
10 (58)			2.43
15 (59)			2.42
20 (60)			1.39
25 (61)			2.13
30 (62)			1.95


- 101 -

(63)			2.61
5 (64)			2.59
10 (65)			1.59
15 (66)			2.31
20 (67)			2.29
25 (68)			2.21


- 102 -




2.20




2.94




2.93





2.96




2.97




1.61




5




(76)




(77)




15




(79)




(80)




(81)




1.63




2.28




2.30




1.66



2.32

















2.34

















1.67

- 103 -


- 104 -

(82)			1.67	
5	(83)			2.35
10	(84)			1.90
15	(85)			2.56
20	(86)			2.58
25	(87)			2.76
30	(88)			2.77

- 105 -


5	(89)			1.88
10	(90)			2.55
15	(91)			2.57
20	(92)			1.74
25	(93)			2.39
30	(94)			2.39
	(95)			1.39

- 106 -




5

2.11




10

2.11




15

2.44




20

2.41



25

2.47



30


2.44

- 107 -



5

2.46




10

2.77




15

2.75





20

2.17



25

1.95











30







1.55

1.47


- 108 -

(109)		1.96
5	(110) 	1.51
10	(111) 	1.43
15	(112) 	2.14
20	(113) 	2.17
25	(114) 	2.37
30	(115) 	2.05
	(116) 	2.03

- 109 -


5	(117) 	2.67
10	(118) 	1.60
15	(119) 	2.29
20	(120) 	2.39
25	(121) 	2.11
30	(122) 	2.11

- 110 -




2.62

5




2.61

10




2.47

15




2.49

20




2.52

25




2.52


30



5




10




2.62

15




2


20




25




30




2.47




2.48




2.62



2.62




2.09




2.81

- 111 -



- 112 -



- 113 -





- 114 -



1.69

1.57



5



1.69

1.57



10



2.40

1.70



15



2.43

1.81



20



1.67

1.78

25











2.37













1.97

30


- 116 -

			1.94
5	(160)		1.57
10	(161)		1.91
15	(162)		1.63
20	(163)		1.62
25	(164)		1.93
30	(165)		2.05


- 117 -

	(166)			1.87
5	(167)			1.72
10	(168)			1.45
15	(169)			1.74
20	(170)			1.74
25	(171)			1.79
30				


- 118 -




5




10




15



20




25




30


- 119 -




5




10




15




20





25





30





- 120 -




1.94





1.97





1.82



1.81





1.79





1.81



- 121 -





1.83





1.79





1.98



1.97




2.09



2.05

- 122 -




2.06

5




1.96

10




2.11

15




2.85

20



1.97


25



1.95


30

- 123 -




1.64

5




1.67

10




1.43

15




1.49

20




2.19

25




2.20


- 124 -




1.93




1.93



2.61




2.63




2.01


- 125 -




1.99




2.63




2.65



2.15




2.12




5

2.76




10

2.80




15

1.99




20

1.96



25

2.65




5

2.64


The present invention further relates to compounds (225) to (449) of formula A-NH-CO-NH-(CHMe)-B, wherein Me is a methyl group and A and B are as given in the table below:

	A	B	Rt
10			
15	(225)		
20	(226)		
25	(227)		
30	(228)		
	(229)		

- 128 -



- 129 -



- 130 -




5




10




15




20




25




30



5



10



15



20









25




30


- 131 -




- 132 -



- 133 -




- 134 -




5




10




15




(275)



20




25




30


- 135 -




5




10




15




(282)



20




25




30

- 136 -









- 137 -










- 138 -




5





30




- 139 -




5




10









15



20




25




30

- 140 -




30.

- 141 -




30


- 142 -




5




10




15



20




25




30


- 143 -




5




10




15






20




25




30




- 144 -




- 146 -




5




10




15









20




25



30




- 147 -




2.62

5




2


10



15



20





25



2.86

30



- 148 -





5




10





15



20





25





30



- 149 -





5





10





15



20




25




30


- 150 -




5




10




15









20




25




30




- 151 -




5




10




15









20




25




30




- 152 -



- 153 -




- 154 -




(397)

5




(398)

10




(399)

15




(400)

20



(401)


25



(402)


30

- 155 -




(403)

5




(404)

10




(405)

15




(406)

20









(407)

25















(408)

30















- 156 -




30

- 157 -




30


- 158 -




5




10




15



20




25




30


- 159 -




5




10




15









20













25













30




- 160 -



- 161 -




- 162 -




5


2.81




10




15



20




25



30

2.69

- 163 -



5

The present invention further relates to compounds (449) to (672) of formula A-NH-CO-CR⁶R⁷-NH-B, wherein R⁶ and R⁷ form, together with the carbon atom of the methylene moiety they are bound to, a cyclopropane moiety, and wherein A and B are as given in the table below:


10

A		B	Rt
(449)			2.13
(450)			2.11
(451)			
(452)			


25

30


- 164 -




- 165 -




- 166 -




5




10




15









20




25




30




- 167 -




5




10




15



20




25




30


- 168 -




- 169 -




- 170 -




5




10




15




20



(498)




25




30


- 171 -




5




10




15




20



(505)




25




30


- 172 -




- 173 -




- 174 -




5




10





15



20




25




30


- 175 -




5





10




15



20






25









30

- 176 -




30

- 177 -




30


- 178 -




5




10




15









20




25




30




- 179 -




5




10



15




20











25




(558) 




30




- 180 -




- 181 -




- 182 -




5




10




15



20




25




30


- 183 -




5




10




15









20




25




30




- 184 -




5




10




15



20




25




30


- 185 -




5




10




15




20




25




30




5




10




15










20




25




30




5




10




15




20














25



30















- 188 -




30

- 189 -




30


- 190 -




5




10




15




20




25




30




5




10




15




20










25





30





- 191 -





- 192 -





5





10





15



20





25





30



- 193 -





5





10





15




20




25




30




5




10




15










20




25




30




5




10




15









20



25




30




- 196 -


(659)




5





(660)



10





15 (661)




20

(662)




25

(663)




30




- 197 -


(664)




5





(665)



10





15 (666)




20

(667)





25

(668)




30






5




(670)

10




(671)

15



(672)


20



25 The present invention further relates to compounds (673) to (758) as given in the table below:


Compound	Rt
<p>(673)</p>	2.32

30




2.20

5




10

2.99




15

2.93




20

2.68



25

2.44



30

2.40


- 200 -

5	(680)		2.77
10	(681)		2.55
15	(682)		2.25
20	(683)		2.86
25	(684)		3.20
30	(685)		3.01


- 201 -

5	(686)		2.87
10	(687)		3.19
15	(688)		4.52 ^a
20	(689)		4.78 ^a
25	(690)		4.41 ^a
30	(691)		4.97 ^a


- 202 -

4.59^a


5

5.28^a


10

4.42^a


15

3.06^b


20

3.02^b


25

1.30^b


- 203 -

3.49^a


5

3.94^a


10

4.31^a


15

2.44^b

20

5.39^a

25



2.91

30


- 204 -

5	(704)		4.59 ^a
10	(705)		4.61 ^a
15	(706)		4.71 ^c
20	(707)		2.31 ^b
25	(708)		4.85 ^c
30	(709)		2.33 ^b


- 205 -

5	(710)		2.66
10	(711)		3.03 ^b
15	(712)		4.68 ^b
20	(713)		4.47 ^a
25	(714)		3.29 ^a
30	(715)		4.03 ^a

- 206 -




5

4.35^a


10

2.41



15

2.13




25


2.51^b

30

- 207 -




5

2.09^b


10

2.55




15

2.95



25

2.68



30

2.19


- 208 -

5	(725)		3.42 ^b
10	(726)		2.83
15	(727)		2.61
20	(728)		3.19 ^b
25	(729)		3.38 ^b
30	(730)		3.59 ^a


- 209 -

5	(731)		4.91 ^a
10	(732)		3.59 ^a
15	(733)		2.17
20	(734)		3.11 ^b
25	(735)		3.29 ^b
30	(736)		3.21 ^b


- 210 -



5


2.51^b

10

2.79^b


15

2.60




20

2.47



25


2.69




30

3.41^b


- 211 -




5

3.23^b


10

3.39^b

15

3.23^b

20





2.64^b

25

3.31^b






30

- 212 -




30

- 213 -




30

- 214 -



5




10


15 The present invention further relates to compounds (759) to (825) as given in the table below:

Compound	Rt
20	
<p>(759)</p>	2.03
25	
<p>(760)</p>	1.99
30	


- 215 -




5




10



15








20







25

30


- 216 -




- 217 -




- 218 -




2.2



2.19




2.18




2.23

30


- 219 -




2.66




2.09



2.13







(782)

3.01^a

30


- 220 -

2.95^a2.07^a3.01^a


2.11

30


- 221 -




2.03




1.93



2.08







2.05







1.97


- 222 -




- 223 -



- 228 -







- 229 -




- 230 -

The present invention further relates to compounds (826) to (874) as given in the table below:


Compound	Rt
 (826)	5
 (827)	10
 (828)	15
 (829)	20

30


- 231 -




5




10



15



20




25


2.67

30


- 232 -



2.83



2.69




20


25

30


- 233 -



2.61



2.84




2.88

20

25


30

- 234 -




5

(841)




10

(842)




20

(843)




25

(844)




5

(845)




10

(846)



15





(847)



20





(848)

- 236 -




30

- 237 -




30

- 238 -




5

(857)



10


(858)



20


(859)

- 239 -




5

(860)




10

(861)



20







(862)




25


(863)

30






(873)




(874)

The nomenclature as used herein for defining compounds, especially the compounds according to the invention, is in general based on the rules of the IUPAC-organisation for chemical compounds and especially organic compounds.


In a special embodiment, one or more of the methylene urea derivatives according to sub formulae Ia to Ix and/or compounds (1) to (224), compounds (225) to (448), compounds (449) to (672), compounds (673) to (758), compounds (759) to (825) and/or compounds (826) to (874) additionally comprise one or two substituents selected from the group consisting of $O(CH_2)_nNR^{11}R^{12}$, $NR^{11}(CH_2)_nNR^{11}R^{12}$, $O(CH_2)_nOR^{12}$, $NR^{11}(CH_2)_nOR^{12}$, wherein



(870)



(871)




(872)

- 244 -


R^{11}, R^{12} are independently selected from a group consisting of H, A, $(CH_2)_mAr^3$ and $(CH_2)_mHet$, or in $NR^{11}R^{12}$,

- R^{11} and R^{12} form, together with the N-Atom they are bound to, a 5-, 6- or 7-membered heterocycly which optionally contains 1 or 2 additional hetero atoms, selected from N, O and S, and n is 1, 2, 3, 4, 5 or 6, preferably 2, 3 or 4.

- In this special embodiment, the substituents are preferably selected from the group consisting of $HNC_2H_4NH_2$, $OCH_2CH_2NH_2$, $NHCH_2CH_2OH$, $OCH_2CH_2NHCH_3$, $N(CH_3)CH_2CH_2NH_2$, $HN(CH_3)CH_2CH_2NH$, $N(CH_3)CH_2CH_2N(CH_3)_2$, $N(CH_3)CH_2CH_2N(CH_3)_2$, $N(CH_3)CH_2CH_2OCH_3$, $OCH_2CH_2N(CH_3)_2$, $OCH_2CH_2N(CH_2CH_3)_2$ and compounds of the formulae



and/or compounds of formulae



- In a further special embodiment, one or more of the methylene urea derivatives according to sub formulae IIa to IIx and/or compounds (1) to

- 245 -

(224), compounds (225) to (448), compounds (449) to (672), compounds (673) to (758), compounds (759) to (825) and/or compounds (826) to (874) additionally comprise one or two substituents selected from the group consisting of $(CH_2)_nS(O)_uNR^{11}R^{12}$ and $(CH_2)_nS(O)_uR^3$ wherein R^{11} , R^{12} and R^3 are defined as above and n is as defined above, preferably n is 0, 1 or 2 and especially is 0, and u is preferably 2 or 3. In this embodiment, the residues are preferably selected from SO_2CH_3 , SO_2CF_3 , OSO_2CH_3 , OSO_2CF_3 , SO_2NH_2 , $SO_2NHCH(CH_3)_2$, $SO_2N(CH_3)_2$, $SO_2N(CH_2CH_3)_2$ and 4-Morpholino-sulfonyl.

In this special embodiment, the additional substituents are preferably bound to one of the aromatic residues directly bound to the methylene urea moiety and/or the pyridinyl residue. More preferably, one or two additional substituents are bound to the residue Ar^1 according to formula II. Even more preferably, in one or more of the formulae IIa to IId, one or two additional substituents are bound to the phenyl moiety directly bound to the nitrogen atom of the methylene urea moiety, i. e. the phenyl moiety at the left hand side of the respective formulae. Especially preferred are compounds (1) to (224), compounds (225) to (448) and/or compounds (449) to (672), wherein one or two additional substituents are bound to the moiety A.

Another aspect of the invention relates to a method for producing compounds of formula II, characterised in that

- a) A compound of formula III



wherein


EG is a functional group, selected from

 $-\text{N}=\text{C}=\text{Y}$ and $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$,

wherein Y is as defined above and below, LG is a leaving group, preferably a leaving group selected from OR^{25} and CHal_3 , wherein R^{25} is selected from the group consisting of unsubstituted or substituted aromatic residues, unsubstituted or substituted heteroaromatic residues and $(\text{O})_2\text{S-R}^{26}$, wherein R^{26} is selected from unsubstituted or substituted aromatic residues and unsubstituted or substituted alkyl residues residues, and wherein R^p , p and Ar^1 are as defined above and below.

is reacted

b) with a compound of formula IV,



wherein

L^2, L^3 are independently from one another H or a metal ion, and $R^6, R^7, E, G, M, Q, U, R^9, q, X, A^2, R^{10}$ and r are as defined above and below,

and optionally

c) isolating and/or treating the compound of formula II obtained by said reaction with an acid, to obtain the salt thereof.

The compounds of the formula I and preferably the compounds of the formula II and also the starting materials for their preparation are, in addition, prepared by methods known per se, as described in the literature (for example in the standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart), to be precise under reaction conditions which are known and suitable for the said reactions. Use can also be made here of variants which are known per se, but are not mentioned here in greater detail.

10 If desired, the starting materials can also be formed in-situ by not isolating them from the reaction mixture, but instead immediately converting them further into the compounds of the formula I or II, respectively. On the other hand, it is possible to carry out the reaction stepwise.

15 The compounds of the formula I and especially the compounds of formula II can preferably be obtained by reacting compounds of the formula III with compounds of the formula IV.

In compounds of formula III, the group FG is a suitable functional group that this preferably selected from -N=C=Y and -NH-(C=Y)-LG . In the functional groups -N=C=Y and/or -NH-(C=Y)-LG , Y is preferably selected from the group consisting of O, S, NR^{21} , $\text{C(R}^{22}\text{)-NO}_2$, $\text{C(R}^{22}\text{)-CN}$ and C(CN)_2 , and more preferably selected from O, S and NR^{21} , even more preferably selected from O and S and especially is O, wherein R^{21} and R^{22} are as defined above/below.

In the compounds of formula III, wherein FG is $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$, LG is a suitable leaving group. Suitable leaving groups are known in the art, for example from Houben-Weyl, Methods of Organic chemistry. Preferably, the leaving group is selected from CHal_3 , wherein Hal is as defined above/below and preferably is chlorine or bromine and especially is chlorine, and OR^{25} , wherein R^{25} is selected from the group consisting of unsubstituted or substituted aromatic residues, unsubstituted or

- 248 -

substituted heteroaromatic residues and $(O)_2S-R^{28}$, wherein R^{28} is selected from unsubstituted or substituted aromatic residues and unsubstituted or substituted alkyl residues residues, and wherein R^8 , p and Ar^1 are as defined above and below.

5 In compounds of formula III in which FG is $-NH(C=Y)LG$ and LG is OR^{25} , $CHal_3$, Hal is preferably selected independently from one another from the group consisting of chlorine, bromine and iodine, even more preferably chlorine and bromine and especially preferred chlorine. Preferably, $CHal_3$ is selected from the group consisting of CCl_3 and CBr_3 and especially preferred $CHal_3$ is CCl_3 .

10 In compounds of formula III in which FG is $-NH(C=Y)LG$ and LG is OR^{25} , R^{25} is preferably selected from unsubstituted or substituted phenyl moieties, preferably substituted phenyl moieties which comprises one or more nitro groups ($-NO_2$) and/or one or more sulfonic acid groups ($-SO_3H$) or salts thereof as substituents, and $(O)_2S-R^{28}$, wherein R^{28} is selected from unsubstituted or substituted phenyl moieties, preferably alkyl substituted phenyl moieties, and unsubstituted or substituted alkyl residues residues, preferably unsubstituted or substituted C_1-C_4 -alkyl moieties and especially unsubstituted or substituted methyl moieties. Substituted alkyl moieties preferably comprise one or more halogen substituents up to pentafluoro. Preferred as halogen substituents are fluorine and chlorine and especially preferred is chlorine. Especially preferred as substituted alkyl moiety is $-CF_3$. Examples of preferred leaving groups OR^{25} are the para-Tosyl- (i. e. $p-Me-C_6H_4-SO_2-$) group, the para-Nitro-phenolate-group (i.e. the $p-O_2N-C_6H_4-O-$) group and the triflate- (i. e. the F_3C-SO_3-) group.

15 If compounds of formula II are desired wherein Y is other than O, it can be advantageous however to carry out the reaction of a compound of formula III, wherein Y is O, and a compound of formula IV according to the invention to obtain a compound of formula II, wherein Y is O, and to modify

- 249 -

or convert the corresponding C=O group (i. e. the C=Y group, wherein Y is O) in the compound of formula II into a $C=NR^{21}$, $C=C(R^{22})NO_2$, $C=C(R^{22})CN$ or $C=C(CN)_2$ group according to methods known in the art, for example from Houben-Weyl, Methods of Organic Chemistry.

5 In the compounds of formula IV, L^2 and/or L^3 is preferably H or a moiety which activates the amino group it is bonded to, for example a metal ion. Suitable metal ions are preferably selected from the group consisting of alkaline metal ions, alkaline-earth metal ions and aluminium ions.

10 Especially preferred metal ions are alkaline metal ions, of which Li, Na K are especially preferred. In case of multi-valent metal ions, the metal ions and the compounds of formula IV form a complex containing one or more compounds of formula IV and one or more metal ions wherein the ratio between compounds of formula IV and metal ions is depending on the valency of the metal ion(s) according to the rules of stoichiometry and/or electroneutrality. Preferably, L^2 or L^3 and more preferred L^2 and L^3 are hydrogen.

15 In detail, the reaction of the compounds of the formula III with the compounds of the formula IV is carried out in the presence or absence of a preferably inert solvent at temperatures between about -20°C and about 200°C , preferably between 0°C and 150°C and especially between room temperature (25°) and 120° . In many cases, it is advantageous to combine one compound of formula III with one compound of formula IV at the lower end of the given temperature range, preferably between -20°C and 75°C , and more preferred between 0°C and 60°C and especially between 10°C and 40°C , for example at about room temperature, and heat the mixture up to a temperature at the upper end of the given temperature range, preferably between 65°C and 180°C , more preferred between 75°C and 150°C and especially between 80°C and 120°C , for example at about 80°C , at about 90°C or at about 100°C . Proceeding in that manner can be advantageous especially in the case that FG is selected from $-NH(C=Y)-$

LG. If RG is selected from $-N=C=Y$ and preferably is $-N=C=O$ or $-N=C=S$ and especially is $-N=C=O$, the reaction can be regularly carried out without prolonged heating to higher temperatures. For example it can be carried out at a temperature between 0 °C and 60 °C and preferably at about room temperature.

5

The reaction between the compounds of formula III, wherein FG is $-NH-(C=Y)-LG$ and especially wherein LG is $CHal$, and compounds of formula IV is preferably carried out in the presence of an acid binding means, for example one or more bases. Suitable acid binding means are known in the art. Preferred as acid binding means are inorganic bases and especially organic bases. Examples for inorganic bases are alkaline or alkaline-earth hydroxides, alkaline or alkaline-earth carbonates and alkaline or alkaline-earth bicarbonates or other salts of a weak acid and alkaline or alkaline-earth metals, preferably of potassium, sodium, calcium or cesium. Examples for organic bases are triethyl amine, diisopropyl ethyl amine (DIPEA), diaza bicyclo undecen (DBU), dimethyl aniline, pyridine or choline. If an organic base is used, it is advantageous in general to use a base with a boiling point that is higher than the highest reaction temperature employed during the reaction. Especially preferred as organic bases are DBU and DIPEA. DBU is especially preferred in the case that LG is $CHal$. DIPEA is especially preferred in the case that LG is OR^{25} .

15

20

Reaction times are generally in the range between some minutes and several days, depending on the reactivity of the respective compounds and the respective reaction conditions. Suitable reaction times are readily determinable by methods known in the art, for example reaction monitoring. Based on the reaction temperatures given above, suitable reaction times generally lie in the range 10 min and 36 hrs, preferably 30 min and 24 hrs and especially between 45 min and 16 hrs, for example about 1 h, about 2 hrs, about 4 hrs, about 6 or about 16 hrs.

30

Preferably, the reaction of the compounds of the formula III with the compounds of the formula IV is carried out in the presence of a suitable solvent, that is preferably inert under the respective reaction conditions. Examples of suitable solvents are hydrocarbons, such as hexane,

5

petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichlorethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; alcohols, such as methanol, ethanol, isopropanol, n-propanol, n-butanol or tert-butanol; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); ketones, such as acetone or butanone; amides, such as acetamide, dimethylacetamide, dimethylformamide (DMF) or N-methyl pyrrolidinone (NMP); nitriles, such as acetonitrile; sulfoxides, such as dimethyl sulfoxide (DMSO); nitro compounds, such as nitromethane or nitrobenzene; esters, such as ethyl acetate, or mixtures of the said solvents. Polar solvents are in general preferred. Examples for suitable polar solvents are chlorinated hydrocarbons, alcohols, glycol ethers, nitriles, amides and sulfoxides or mixtures thereof. More preferred are chlorinated hydrocarbons, especially dichloromethane, and sulfoxides, especially DMSO.

10

15

20

Preferably, the reaction between a compound of formula III, wherein FG is $-N=C=Y$ and preferably is $-N=C=O$ or $-N=C=S$ and especially is $-N=C=O$, and a compound of formula IV, especially a compound of formula IV,

25

wherein L^2 and L^3 is H, is carried out in an inert solvent at the lower end of the given temperature range, for example in a chlorinated hydrocarbon, for example dichloromethane, in a temperature range between 0 °C and 60 °C, preferably at about room temperature. Reaction times generally lie in the range of 2 hours to 24 hrs, for example at about 16 hrs. Preferably, no acid binding means is present.

30

- 252 -

Preferably, the reaction between a compound of formula III, wherein FG is $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$ and especially wherein LG is CHal_3 , and compounds of formula IV, especially a compound of formula IV, wherein L^2 and L^3 is H, is carried out in an inert solvent, preferably a solvent boiling at higher temperatures, for example a sulfoxide and especially DMSO, in a temperature range between 60 °C and 120 °C, for example at about 80 °C. Reaction times generally lie in the range of 1 hrs to 10 hrs, for example between 2 and 6 hrs. Preferably, the reaction is carried out in the presence of an acid binding means, preferably one of the afore mentioned acid binding means, more preferably an organic base and especially in the presence of DBU.

Preferably, the reaction between a compound of formula III, wherein FG is $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$ and especially wherein LG is OR^{25} , and compounds of formula IV, especially a compound of formula IV, wherein L^2 and L^3 is H, is carried out in an inert solvent at the lower end of the given temperature range, for example in a chlorinated hydrocarbon, for example dichloromethane, in a temperature range between 0 °C and 60 °C, preferably at about room temperature. Reaction times generally lie in the range of 2 hours to 24 hrs. Preferably, the reaction is carried out in the presence of an acid binding means, preferably one of the afore mentioned acid binding means, more preferably an organic base and especially in the presence of DIPEA.


In general, the compounds of formula III and/or formula IV are new. In any case, they can be prepared according to methods known in the art.

The compounds of formula III can be obtained according to methods known in the art. In an advantageous manner, they can be readily obtained by one or more of the reaction routes given below:

- 253 -

Compounds of formula III, wherein FG is $-\text{N}=\text{C}=\text{Y}$ and Y is O or S can be readily obtained from suitable substituted derivatives of $(\text{R}^8)_p\text{-Ar}^1$ according to known procedures for producing isocyanates and thiocyanates. When FG is $-\text{N}=\text{C}=\text{O}$, the compounds of formula III can be readily obtained via Curtius-, Hoffmann or Lossen rearrangement starting from $(\text{R}^8)_p\text{-Ar}^1\text{-COOH}$ or the respective acid halides, as described in the art. If desired, compounds of formula III, wherein Y is O can be readily derivatized to compounds of formula III, wherein Y is S, according to procedures known in the art.

Compounds of formula III, wherein FG is $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$ and especially wherein LG is CHal_3 can be readily obtained from the reaction of suitable amino substituted derivatives of $(\text{R}^8)_p\text{-Ar}^1$ of formula V




wherein R^8 , p, and Ar^1 are as defined above/below and L^4 and L^5 are selected independently from each other from the meanings given for L^2 and L^3 and more preferred are hydrogen, with a compound of formula VI



wherein Y is as defined above/below and L^6 is preferably selected from Cl, Br, I, OH, reactive derivatized OH-moieties, especially reactive esterified OH-moieties, for example alkylsulfonyloxy-moieties comprising 1 to 6 carbon atoms (preferably methylsulfonyloxy) or arylsulfonyloxy-moieties comprising 6 to 10 carbon atoms (preferably phenyl- or aryl-p-

tolylsulfonyloxy), and diazonium moieties, and more preferred selected from Cl, Br or I, and even more preferred is Cl.

Compounds of formula III, wherein FG is -NH-(C=Y)-LG and especially wherein LG is CHal_3 can be readily obtained from the reaction of suitable amino substituted derivatives of $(\text{R}^6)_p\text{-Ar}^1$ of formula V



wherein R^6 , p, and Ar^1 are as defined above/below and L^4 and L^5 are selected independently from each other from the meanings given for L^2 and L^3 and more preferred are hydrogen, with a compound of formula VIa



wherein Y and L^6 are as defined above/below.

The reaction between compounds of formula V and compounds of formula VI can be carried out in the presence of a suitable solvent, that is preferably inert at the chosen reaction conditions. Suitable solvents are known in the art. Examples of suitable solvents include hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; nitriles, such as acetonitrile; esters, such as ethyl acetate, or mixtures of said solvents. Non-protic solvents are in general preferred.


The reaction between compounds of formula V and compounds of formula VI can be carried out in the presence of a suitable acid binding means, especially organic or anorganic bases. Examples for inorganic bases are alkaline or alkaline-earth hydroxides, alkaline or alkaline-earth carbonates and alkaline or alkaline-earth bicarbonates or other salts of a weak acid and alkaline or alkaline-earth metals, preferably of potassium, sodium, calcium or cesium. Examples for organic bases are triethyl amine, diisopropyl ethal amine (DIPEA), diaza bicyclo undecan (DBU), dimethyl aniline, pyridine or chinoline. If an organic base is used, it is advantageous in general to use a base with a boiling point that is higher than the highest reaction temperature employed during the reaction.

The reaction between compounds of formula V and compounds of formula VI can be carried out in the presence of a suitable solvent, that is preferably inert at the chosen reaction conditions. Suitable solvents are known in the art. Examples of suitable solvents include hydrocarbons, such as hexane, petroleum ether, benzene, toluene or xylene; chlorinated hydrocarbons, such as trichloroethylene, 1,2-dichloroethane, tetrachloromethane, chloroform or dichloromethane; ethers, such as diethyl ether, diisopropyl ether, tetrahydrofuran (THF) or dioxane; nitriles, such as acetonitrile; esters, such as ethyl acetate, or mixtures of said solvents. Non-protic solvents are in general preferred.

If the reaction between a compound of formula V and a compound of formula VI is carried out in presence of an organic base that is liquid at the chosen reaction conditions, it can be advantageous not to add an additional solvent.

Compounds of formula III, wherein FG is -NH-(C=Y)-LG and preferably wherein LG is OR^{25} and especially wherein R^{25} is an unsubstituted or substituted phenyl moiety, can be readily obtained from the reaction of suitable amino substituted derivatives of $(\text{R}^6)_p\text{-Ar}^1$ of formula V

- 256 -



V

wherein R^8 , p, and Ar^1 are as defined above/below and L^4 and L^5 are selected independently from each other from the meanings given for L^2 and L^3 and more preferred are hydrogen, with a compound of formula VIb



VIb

wherein Y and L^6 are as defined above/below.

15 Suitable reaction conditions for carrying out reaction of compounds of formula V with compounds of formula VI, VIa and VIb, respectively, are known in the art. In detail, the reaction of the compounds of the formula V with the compounds of the formula VI is carried out in the presence or absence, preferably in the presence of an inert solvent, preferably one of the afore mentioned inert solvents, more preferably ethers and chlorinated hydrocarbons, and especially in dichloromethane, preferably in a temperature range between 0 °C and 60 °C and more preferably at about room temperature.

25 The reaction between compounds of formula V and compounds of formula VI is preferably carried out in the presence of an acid binding means, for example one or more bases. Suitable acid binding means are known in the art. Preferred as acid binding means are organic bases, more preferably one of the afore mentioned organic bases and especially pyridine.


- 257 -

In general, the reaction times for the reaction between compounds of formula V and compounds of formula VI lie in the range between 6 hrs and 36 hrs, preferably 12 hrs to 24hrs, for example at about 16 hrs.

5 Some of the starting materials of the formula V and/or the formula VI are known and preferably commercially available. If they are not known, they can be prepared by methods known per se.


10 The compounds of formula IV can be obtained according to methods known in the art.

If the compound of formula IV is a compound according to formula IVa,




IVa

15 it can be readily obtained in an advantageous manner (reaction route A) by reacting a compound of formula VIIa,



VIIa

25 wherein E, G, M, Q, U, R^9 and q are as defined above/below, with a compound of formula VIII,




VIII

wherein L^8 is H or a metal ion, preferably a metal ion selected from the group consisting of alkaline metal ions, alkaline-earth metal ions and aluminum ions, especially preferred alkaline metal ions, of which Li, Na and K are especially preferred, and even more preferred is H; and Ar^2 , R^{10} , r and X are as defined above/below, and preferably wherein X is $(CHR^{11})_h$ - $Q-(CHR^{12})_h$, wherein R^{11} , h and i and R^{12} are defined above/below, and more preferred wherein h and/or i is 0 and Q is selected from a group consisting of O, S, $N-R^{15}$, $(O-CHR^{15}O)_j$, $(O-CHR^{15}CHR^{19})_j$, $O-N=CH$, $NR^{15}-N=CH$, $NR^{15}SO_2$, wherein R^{15} , R^{18} , R^{19} and j are as defined above/below, and even more preferred wherein h and i is 0 and Q is selected from a group consisting of O, S, $N-R^{15}$;

optionally isolating the reaction product,


and transferring the obtained reaction product of formula IX




into a compound of formula IVa, preferably by reducing and more preferred by hydrogenating the CN-moiety of the compound of formula IX into a H_2NCH_2 -moiety. Methods and reaction conditions for hydrogenating said CN-moiety into a H_2NCH_2 -moiety are known in the art. In general, it is advantageous to carry out the hydrogenation reaction in the presence of a hydrogen delivering means, for example hydrogen gas, in the presence of a suitable catalyst, preferably a Nickel catalyst, for example Raney-Nickel. In general, such hydrogenation reactions are carried out in a suitable solvent. Suitable solvents for hydrogenation reactions are known in the art. Suitable solvents, for example, are alcohols, especially methanol and ethanol and ethers, especially THF, and mixtures thereof. Preferably, the

hydrogenation reaction is carried out in a methanol/ammonia mixture, preferably in the presence of Raney nickel. In general, the hydrogenation reactions are carried out at about normal pressure or elevated pressure, for example between normal pressure and 10 bar pressure, preferably at about 5 bar pressure (about 500 kPa). The hydrogenation reaction is usually carried out in the temperature range between $-20^\circ C$ and $150^\circ C$, preferably $+20^\circ C$ and $100^\circ C$, for example at about $45^\circ C$.

Ar^2 is preferably pyridinyl. Accordingly, the compound of formula VIII is preferably selected from the group consisting of formulae VIIa and VIIb,




VIIa




VIIb

wherein L^8 , X, R^{10} and r are as defined above, and especially preferred from the group consisting of formulae VIIc and VIId,



VIIc



VIId


wherein R^{10} and r are as defined above, or the alkaline metal salts and especially the sodium or potassium salts thereof.

Accordingly, in formulae IVa, VII, VIIa, VIIb and IX, the bridging group X is preferably O, S, OCH_2 and OCH_2CH_2 and especially is O.

- 260 -


In the formulae VII, VIIa and VIIb, L⁸ is preferably H or selected from the group consisting of Na, K and Cs and especially preferred is H.

5 In general, this reaction is advantageous to produce compounds of formula IVaa,



wherein E, G, M, U, R⁹, q, X, Ar², R¹⁰ and r are as defined above/below.


15 To obtain compounds of formula IVaa, it is reasonable to employ a compound of formula VII that is selected from the compounds of formula VIIa,



and proceed the reaction as described above/below.


25 Accordingly, by starting from a compound of formula VIIa and a compound of formula VIIa, the reaction preferably leads to compounds of formula IVaaa,

- 261 -




wherein E, G, M, U, R⁹, q, X, R¹⁰ and r are as defined above/below.

10 Accordingly, by starting from a compound of formula VIIa and a compound of formula VIIb, the reaction preferably leads to compounds of formula IVaab,




wherein E, G, M, U, R⁹, q, X, R¹⁰ and r are as defined above/below.

20 Accordingly, by starting from a compound of formula VIIa and a compound of formula VIIIc, the reaction preferably leads to compounds of formula IVaac,



wherein E, G, M, U, R⁹, q, R¹⁰ and r are as defined above/below.

30 Accordingly, by starting from a compound of formula VIIa and a compound of formula VIIIc, the reaction preferably leads to compounds of formula



5 wherein E, G, M, U, R⁹, q, R¹⁰ and i are as defined above/below.

Some of the starting materials of the formula VII and/or the formula VIII are known and preferably commercially available. If they are not known, they can be prepared by methods known per se.

10

The reaction between the compound of formula VII and VIII is preferably carried out in the temperature range between 0 °C and 250 °C, more preferred room temperature and 200 °C, for example at about 120 °C, at about 150 °C or at about 180°. Reaction times depend on the respective reactants and the respective reaction temperature, but generally lie in the range between 30 min and 36 hrs, preferably 3 hrs and 24 hrs, more preferably 8 hrs and 20 hrs for example about 10 hrs, about 16 hrs or about 18 hrs.

15

20 The reaction can be carried out in the absence of solvent or preferably in the presence of a solvent, preferable a solvent that is inert under the respective reaction conditions. Suitable inert solvents for carrying out the reaction are known in the art. Examples for suitable solvents are high boiling aliphatic hydrocarbons, high boiling aromatic carbons, for example toluene, xylenes, high boiling chlorinated hydrocarbons, such as trichloroethylene, tetrachloroethanes, pentachloroethanes and hexachloroethanes; high boiling ethers, such as ethylene glycol and propylene glycols; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); amides, such as acetamide, dimethylacetamide, dimethylformamide (DMF) or N-methyl pyrrolidinone (NMP); sulfoxides, such as dimethyl sulfoxide (DMSO); or


30

mixtures of the said solvents. Preferred are amides, especially dimethylformamide (DMF) or N-methyl pyrrolidinone (NMP).

5 Preferably, the reaction is carried out in the presence of a base. Suitable bases are known in the art. Preferred bases are organic bases and especially inorganic bases. Examples for inorganic bases are alkaline or alkaline-earth hydroxides, alkaline or alkaline-earth carbonates and alkaline or alkaline-earth bicarbonates or other salts of a weak acid and alkaline or alkaline-earth metals, preferably of potassium, sodium, calcium or cesium. Preferred inorganic bases are K₂CO₃, Na₂CO₃, MgCO₃, CaCO₃, NaOH and KOH, especially preferred is K₂CO₃. Examples for organic bases are triethyl amine, diisopropyl ethyl amine (DIPEA), dimethyl aniline, pyridine or chinoline. If an organic base is used, it is advantageous in general to use a base with a boiling point that is higher than the highest reaction temperature employed during the reaction.

15


Alternatively, if the compound of formula IV is a compound according to formula IVb,



20

it can be readily obtained in an advantageous manner (reaction route B) by reacting a compound of formula VIIb,

25




30

- 264 -

- wherein E, G, M, Q, U, R⁶ and q are as defined above/below and wherein L⁹ is selected independently from H or a moiety which activates the group (and preferably a hetero atom such as N, S and especially O which is part of the group) it is bonded to, for example a metal ion. Suitable metal ions are preferably selected from the group consisting of alkaline metal ions, alkaline-earth metal ions and aluminium ions. More preferred, L⁹ is selected from H, Na and K, and is even more preferred H, especially if X is selected from the group consisting of wherein X is (CHR¹¹)_n-Q-(CHR¹²)_n, wherein R¹¹, h and l and R¹² are defined above/below, and more preferred wherein h and/or l is O and Q is selected from a group consisting of O, S, N-R¹⁵, (CHR¹⁸-O), (CHR¹⁸CHR¹⁹-O), CH=N-O, CH=N-NR¹⁵, SO₂NR¹⁵, wherein R¹⁵, R¹⁸, R¹⁹ and j are as defined above/below, and even more preferred wherein h and l is O and Q is selected from a group consisting of O, S, N-R¹⁵,


- 15 with a compound of formula VIIIb,



VIIIb

- 20 wherein L¹⁰ is preferably Cl, Br, I or diazonium moiety, more preferred Cl, Br or I and even more preferred Br and Cl; optionally isolating the reaction product,

- 25 and transferring the obtained reaction product of formula IXb




30


- 265 -

into a compound of formula IVa, preferably by reducing and more preferred by hydrogenating the CN-moiety of the compound of formula IXa into a H₂NCH₂-moiety, preferably as described above for the compound IX.

- 5 Ar² is preferably pyridinyl. Accordingly, the compound of formula VIIIb is preferably selected from the group consisting of formulae VIIIe and VIIIf,




VIIIe




VIIIf

wherein L¹⁰, R⁹ and r are as defined above, and especially preferred from the group consisting of formulae VIIIg and VIIIh,



VIIIg




VIIIh

- 20 wherein Hal, R⁹ and r are as defined above, and wherein Hal is preferably Cl in compounds of formula VIIIg and preferably Br in compounds of formula VIIIh.

- 25 Accordingly, in formulae IVb, VIIIb, VIIIe, VIIIf and IXb, the bridging group X is preferably O, S, OCH₂ and OCH₂CH₂ and especially is O.


In general, this alternative reaction is advantageous to produce compounds of formula IVbb.

30




wherein E, G, Q, U, R⁹, q, X, A², R¹⁰ and r are as defined above/below.

To obtain compounds of formula IVbb, it is reasonable to employ a compound of formula VIIb that is selected from the compounds of formula VIIbb,



wherein E, G, Q, U, X and L⁹ are as defined above/below, more preferred wherein X-L⁹ is selected from the group consisting of SH, OH and HN-R¹⁷ and especially wherein X-L⁹ is OH, and proceed the alternative reaction as described above/below.


Accordingly, by starting from a compound a formula VIIb and a compound of formula VIIe, the reaction preferably leads to compounds of formula IVbbe,



wherein E, G, Q, U, R⁹, q, X, R¹⁰ and r are as defined above/below.


30

Accordingly, by starting from a compound of formula VIbb and a compound of formula VIIIf, the reaction preferably leads to compounds of formula IVbbf,




wherein E, G, Q, U, R⁹, q, X, R¹⁰ and r are as defined above/below.

Accordingly, by starting from a compound of formula VIbb and a compound of formula VIIg, the reaction preferably leads to compounds of formula IVbbg,



wherein E, G, Q, U, R⁹, q, R¹⁰ and r are as defined above/below.

Accordingly, by starting from a compound of formula VIb and a compound of formula VIIh, the reaction preferably leads to compounds of formula IVbbh,



wherein E, G, Q, U, R⁹, q, R¹⁰ and r are as defined above/below.

- 268 -

Some of the starting materials of the formula VIIb and/or the formula VIIIb are known and preferably commercially available. If they are not known, they can be prepared by methods known per se.

5

The reaction between the compound of formula VIIb and VIIIb is preferably carried out in the temperature range between 0 °C and 250 °C, more preferred 50 °C and 220 °C, for example at about 90 °C, at about 120 °C, at about 160 °C, at about 180 °C or at about 200°. Reaction times depend on the respective reactants and the respective reaction temperature, but generally lie in the range between 10 min and 36 hrs, preferably between 60 min and 24 hrs, more preferably 3 h and 20 hrs for example about 6 hrs, about 12 hrs, about 15 hrs or about 18 hrs.

10

15

The reaction can be carried out in the absence or the presence of a solvent, preferable a solvent that is inert under the respective reaction conditions. Suitable inert solvents for carrying out the reaction are known in the art. Examples for suitable solvents are high aliphatic hydrocarbons, aromatic carbons, for example toluene and xylenes, high boiling chlorinated hydrocarbons, such as dichloromethane, trichloromethane trichloroethylene, tetrachloroethanes, pentachloroethanes and hexachloroethanes; ethers, such as diethylether, tert.-butyl methyl ether, ethylene glycol and propylene glycols; glycol ethers, such as ethylene glycol monomethyl or monoethyl ether or ethylene glycol dimethyl ether (diglyme); nitriles, such as acetonitrile, amides such as acetamide, dimethylacetamide, dimethylformamide (DMF) or N-methyl pyrrolidinone (NMP); sulfoxides, such as dimethyl sulfoxide (DMSO); or mixtures of the said solvents.

25

30

In many cases, it is advantageous to carry out the reaction in the presence of a catalyst. Suitable catalysts are known in the art. Preferred catalysts are compounds comprising catalytically active metals and especially

- 269 -

compounds comprising catalytically active copper. A preferred compound comprising catalytically active copper is copper iodide and especially is CuI. Carrying out the reaction in the presence of a catalyst as described above is preferred if a compound of formula VIII is used, wherein L¹⁰ or Hal is bromine, and is especially preferred if a compound of formula VIIIc or VIIIh is used, wherein L¹⁰ or Hal is bromine.

5

In many cases, it is advantageous to carry out the reaction in the presence of an acid binding means, preferably an organic base as described above and more preferred an inorganic base. Preferred inorganic bases are K₂CO₃, Na₂CO₃, MgCO₃, CaCO₃, NaOH and KOH, especially preferred is K₂CO₃. Carrying out the reaction in the presence of and acid binding means as described above is preferred if a compound of formula VIII is used, wherein L¹⁰ or Hal is bromine, and is especially preferred if a compound of formula VIIIc or VIIIh is used, wherein L¹⁰ or Hal is bromine.

10

15


Preferably, the reaction is carried out by heating up a reaction mixture comprising one compound of formula VIIb and one compound of formula VIIIb to a suitable reaction temperature, which preferably lies at the upper end of the given temperature ranges and more preferred is in the range between 150 °C and 200 °C, for example at about 180 °C, preferably in the presence of the suitable catalyst and especially in the presence of copper. Reaction times at this temperature are preferably as given above and especially in the range between 1 h and 5 hrs, for example about 3 hrs. Preferably, the reaction mixture is then allowed to cool down to a temperature in the lower range of the given temperature, more preferred to a temperature in the range between 50 °C and 150 °C, for example to about 90°. Preferably, a suitable solvent, especially tert.-butyl methyl ether, is then added and the reaction mixture is preferably kept at about the same temperature for some more time, preferably for 30 min to 2 hrs and more preferred for about one hour.

25

30

- 270 -

if the compound IV is a compound according to formula IVc,



5

it can be readily obtained in an advantageous manner (reaction route C) by reacting a compound of formula XI




10

wherein L^9 is H or a metal ion, preferably a metal ion selected from the group consisting of alkaline metal ions, alkaline-earth metal ions and aluminium ions, especially preferred alkaline metal ions, of which Li, Na, and K are especially preferred, and even more preferred H; and E, G, M, Q, U, R^9 , q and X are as defined above/below, and especially wherein X is selected from the group consisting of wherein X is $(CHR^{11})_h-Q-(CHR^{12})_i$, wherein R^{11} , h and i and R^{12} are defined above/below, and more preferred wherein h and/or i is 0 and Q is selected from a group consisting of O, S, $N-R^{15}$, $(CHR^{18}-O)_j$, $(CHR^{18}CHR^{19}-O)_j$, $CH=N-O$, $CH=N-NR^{15}$, SO_2NR^{15} , wherein R^{15} , R^{18} , R^{19} and j are as defined above/below, and even more preferred wherein h and i is 0 and Q is selected from a group consisting of O, S, $N-R^{15}$,

20

25 with a compound of formula XII,




30

- 271 -

wherein hal is independently selected from the group consisting of Cl, Br and I, the residue R^{10} are the same or different and have the meanings given above/below and preferably have both the same meaning, and the indices r are the same or different and have the meanings given above/below and preferably are the same,

5

optionally isolating the reaction product, and transferring the obtained reaction product of formula XIII



10

into a compound of formula IVc, preferably by reducing or hydrogenating the CN-moiety of the compound of formula IX into a H_2NCH_2 -moiety, for example as described above for the compound of formula IX.

15

In the compounds IVc, XII and XIII, r is preferably in each case identical and even more preferred in each case 0.

20

In formulae IVc, XI and XIII, the bridging group X is preferably O, S, OCH_2 and OCH_2CH_2 and especially is O.

25 In the formula XI, L^9 is preferably H or selected from the group consisting of Na, K and Cs and especially preferred is H.

The reaction between the compound of formula XI and XII is preferably carried out in the temperature range between 0 °C and 250 °C, more preferred room temperature and 200 °C, for example at about 120 °C, at about 150 °C or at about 180°. Reaction times depend on the respective reactants and the respective reaction temperature, but generally lie in the

30

- 272 -

range between 30 min and 24 hrs, preferably one hour and 12 hrs, for example about 2 hrs, about 3 hrs or about 6 hrs. The reaction can be carried out in the absence of solvent or in the presence of a solvent, preferably a solvent that is inert under the respective reaction conditions. Suitable inert solvents for carrying out the reaction are known in the art.


In the methods according to the invention for producing compounds, E, G, M, Q, and U are as defined above/below, for example as defined above/below for the compounds according to the invention. More preferably, two or more of E, G, M, Q, and U are carbon atoms. In one embodiment of the method according to the invention for producing compounds, E, G, M, Q, and U all are carbon atoms.

Some of the starting materials of the formula XI and/or the formula XII are known and preferably commercially available. If they are not known, they can be prepared by methods known per se.

Independently of the chosen reaction route, it is in many cases possible or even feasible to introduce residues R^8 , R^9 and/or R^{10} into one or more of the compounds described above, or, if the compound already comprises one or more residues R^8 , R^9 and/or R^{10} , to introduce additional residues R^8 , R^9 and/or R^{10} into said compound. The introduction of additional residues can be readily performed by methods known in the art and especially by aromatic substitution, for example nucleophilic aromatic substitution or electrophilic aromatic substitution. For example, in compounds comprising Ar^1 , wherein Ar^1 comprises one or more halogen and preferably fluorine substituents, one or more of the halogen/fluorine substituents can be easily substituted by hydroxy, thio and/or amino substituted hydrocarbons, preferably selected from the group consisting of $HO(CH_2)_nNR^{11}R^{12}$, $HO(CH_2)_nO(CH_2)_mNR^{11}R^{12}$, $HO(CH_2)_nNR^{11}(CH_2)_kOR^{12}$, $HO(CH_2)_nNR^{11}(CH_2)_kNR^{12}$, $HO(CH_2)_nCOOR^{13}$, $HO(CH_2)_nS(O)_pR^{13}$, $HNR^{11}(CH_2)_nNR^{11}R^{12}$, $HNR^{11}(CH_2)_nO(CH_2)_mNR^{11}R^{12}$,

- 273 -

$HNR^{11}(CH_2)_nNR^{11}(CH_2)_kOR^{12}$, $HNR^{11}(CH_2)_nNR^{11}(CH_2)_kNR^{12}$, $HNR^{11}(CH_2)_nCOOR^{12}$ and $HNR^{11}(CH_2)_nS(O)_pR^{13}$ wherein R^{11} , R^{12} and R^{13} are defined as above and n is as defined above, preferably n is 0, 1 or 2 and especially is 0, k is 1 to 4 and preferably 1 or 2, and u is preferably 2. In this embodiment R^{11} , R^{12} and R^{13} are more preferably selected independently from each other from the group consisting of H, methyl and ethyl. Even more preferred, the hydroxy, thio and/or amino substituted hydrocarbons are selected from the group consisting of NH_2 , $HN(CH_3)_2$, NH_2CH_3 , $HN(C_2H_5)_2$, $H_2NCH_2CH_2NH_2$, $HOCH_2CH_2NH_2$, $HOCH_2CH_2NHCH_3$, $HN(CH_3)CH_2CH_2NH_2$, $HN(CH_3)CH_2CH_2N(CH_3)_2$, $HN(CH_3)CH_2CH_2N(CH_3)_2$, $HN(CH_3)CH_2CH_2OCH_3$, $HOCH_2CH_2N(CH_3)_2$, $HOCH_2CH_2N(CH_2CH_3)_2$, $HSCH_3$, HSC_2H_5 , and compounds of the formulae



or salts and especially metal salts thereof.

On the other hand, it is in many cases possible or even feasible to modify or derivatize one or more of the residue is R^8 , R^9 and R^{10} into residues R^8 , R^9 and/or R^{10} other than the ones originally present. For example, CH_3 groups can be oxidised into aldehyde groups or carbonic acid groups, thio atom containing groups, for example S-alkyl or S-aryl groups, can be oxidised into SO_2 -alkyl or SO_2 -aryl groups, respectively, carbonic acid

groups can be derivatized to carbonic acid ester groups or carbon amide groups and carbonic acid ester groups or carbon amide groups can be hydrolysed into the corresponding carbonic acid groups. Methods for performing such modifications or derivatizations are known in the art, for example from Houben-Weyl, Methods of Organic Chemistry.

5

Every reaction step described herein can optionally be followed by one or more working up procedures and/or isolating procedures. Suitable such procedures are known in the art, for example from standard works, such as Houben-Weyl, Methoden der organischen Chemie [Methods of Organic Chemistry], Georg-Thieme-Verlag, Stuttgart). Examples for such procedures include, but are not limited to evaporating a solvent, distilling, crystallization, fractionised crystallization, extraction procedures, washing procedures, digesting procedures, filtration procedures, chromatography, chromatography by HPLC and drying procedures, especially drying procedures in vacuo and/or elevated temperature.

15

A base of the formula I or the formula II can be converted into the associated acid-addition salt using an acid, for example by reaction of equivalent amounts of the base and the acid in a preferably inert solvent, such as ethanol, followed by evaporation. Suitable acids for this reaction are, in particular, those which give physiologically acceptable salts. Thus, it is possible to use inorganic acids, for example sulfuric acid, sulfurous acid, dithionic acid, nitric acid, hydrohalic acids, such as hydrochloric acid or hydrobromic acid, phosphoric acids, such as, for example,

25

orthophosphoric acid, sulfamic acid, furthermore organic acids, in particular aliphatic, alicyclic, araliphatic, aromatic or heterocyclic monobasic or polybasic carboxylic, sulfonic or sulfuric acids, for example formic acid, acetic acid, propionic acid, hexanoic acid, octanoic acid, decanoic acid, hexadecanoic acid, octadecanoic acid, pivalic acid, diethylacetic acid, malonic acid, succinic acid, pimelic acid, fumaric acid, maleic acid, lactic acid, tartaric acid, malic acid, citric acid, gluconic acid,

30

ascorbic acid, nicotinic acid, isonicotinic acid, methane- or ethanesulfonic acid, ethanesulfonic acid, 2-hydroxyethanesulfonic acid, benzenesulfonic acid, trimethoxybenzoic acid, adamantanecarboxylic acid, p-toluenesulfonic acid, glycolic acid, embonic acid, chlorophenoxyacetic acid, aspartic acid, glutamic acid, proline, glyoxylic acid, palmitic acid, parachlorophenoxyisobutyric acid, cyclohexanecarboxylic acid, glucose 1-phosphate, naphthalenemono- and -disulfonic acids or laurylsulfuric acid. Salts with physiologically unacceptable acids, for example picrates, can be used to isolate and/or purify the compounds of the formula I. On the other hand, compounds of the formula I can be converted into the corresponding metal salts, in particular alkali metal salts or alkaline earth metal salts, or into the corresponding ammonium salts, using bases (for example sodium hydroxide, potassium hydroxide, sodium carbonate or potassium carbonate). Suitable salts are furthermore substituted ammonium salts, for example the dimethyl-, diethyl- and diisopropylammonium salts, monoethanol-, diethanol- and diisopropanolammonium salts, cyclohexyl- and dicyclohexylammonium salts, dibenzylethylenediammonium salts, furthermore, for example, salts with arginine or lysine.

10

15

On the other hand, if desired, the free bases of the formula I or the formula II can be liberated from their salts using bases (for example sodium hydroxide, potassium hydroxide, sodium carbonate or potassium carbonate).

20

The invention relates to compounds of the formula I and of the formula II and physiologically acceptable salts and solvates thereof as medicaments.

25

The invention also relates to the compounds for the formula I and of the formula II and physiologically acceptable salts and solvates thereof as kinase inhibitors.

30

- 276 -

5 The invention furthermore relates to the use of the compounds of the formula I and/or physiologically acceptable salts and/or solvates thereof for the preparation of pharmaceutical compositions and/or pharmaceutical preparations, in particular by non-chemical methods. The invention
10 furthermore relates to the use of the compounds of the formula II and/or physiologically acceptable salts and/or solvates thereof for the preparation of pharmaceutical compositions and/or pharmaceutical preparations, in particular by non-chemical methods. In this cases, one or more compounds according to the invention can be converted into a suitable dosage form together with at least one solid, liquid and/or semi-liquid
15 excipient or adjuvant and, if desired, in combination with one or more further active ingredients.

20 The invention further relates to the use of one or more of the compounds according to the invention, selected from the group consisting of compounds of the formula I as free bases, solvates of compounds of the formula I, salts of compounds of formula I, of compounds of the formula II as free bases, solvates of compounds of the formula II and salts of compounds of formula II, for the production of pharmaceutical compositions and/or pharmaceutical preparations and/or pharmaceutical preparations, in particular by a non-chemical route. In general, non-chemical routes for the production of pharmaceutical compositions and/or pharmaceutical preparations
25 comprise processing steps on suitable mechanical means known in the art that transfer one or more compounds according to the invention into a dosage form suitable for administration to a patient in need of such a treatment. Usually, the transfer of one or more compounds according to the invention into such a dosage form comprises the addition of one or more compounds, selected from the group consisting of carriers, excipients, auxiliaries and pharmaceutical active ingredients other than the compounds according to the invention. Suitable processing steps include, but are not limited to combining, milling, mixing, granulating, dissolving,
30 dispersing, homogenizing, casting and/or compressing the respective

- 277 -

5 active and non-active ingredients. In this respect, active ingredients are preferably at least one compound according to this invention and one or more additional compounds other than the compounds according to the invention, which show valuable pharmaceutical properties, preferably those pharmaceutical active agents other than the compounds according to invention which are disclosed herein.

10 The process for preparing pharmaceutical compositions and/or pharmaceutical preparations preferably comprises one or more processing steps, selected from the group consisting of combining, milling, mixing, granulating, dissolving, dispersing, homogenizing and compressing. The one or more processing steps are preferably performed on one or more of the ingredients which are to form the pharmaceutical composition and/or pharmaceutical preparation preferably according to invention. Even more
15 preferred, said processing steps are performed on two or more of the ingredients which are to form the pharmaceutical composition and/or pharmaceutical preparation, said ingredients comprising one or more compounds according to the invention and, additionally, one or more compounds, preferably selected from the group consisting of active ingredients other than the compounds according to the invention, excipients, auxiliaries, adjuvants and carriers. Mechanical means for performing said processing steps are known in the art, for example from Ullmann's Encyclopedia of Industrial Chemistry, 5th Edition.

20 Preferably, one or more compounds according to the invention are converted into a suitable dosage form together with at least one compound selected from the group consisting of excipients, auxiliaries, adjuvants and carriers, especially solid, liquid and/or semi-liquid excipients, auxiliaries, adjuvants and carriers, and, if desired, in combination with one or more further active ingredients.
30

- 278 -

Suitable dosage forms include, but are not limited to tablets, capsules, semi-solids, suppositories, aerosols, which can be produced according to methods known in the art, for example as described below:

- 5 tablets
- mixing of active ingredient/s and auxiliaries,
compression of said mixture into tablets
(direct compression), optionally granulation
of part of mixture before compression
- 10 capsules
- mixing of active ingredient/s and auxiliaries
to obtain a flowable powder, optionally
granulating powder, filling
powders/granulate into opened capsules,
capping of capsules
- 15 semi-solids (ointments, gels, creams) dissolving/dispersing active
ingredient/s in an aqueous or fatty carrier;
subsequent mixing of aqueous/fatty phase
with complementary fatty resp. aqueous
phase, homogenisation (creams only)
- 20 suppositories (rectal and vaginal) dissolving/dispersing active ingredient/s
in carrier material liquified by heat (rectal:
carrier material normally a wax; vaginal:
carrier normally a heated solution of a
gelling agent), casting said mixture into
suppository forms, annealing and
withdrawal suppositories from the forms
- 25 aerosols:
- dispersing/dissolving active agent/s in a
propellant, bottling said mixture into an
- 30

- 279 -

atomizer

The invention thus relates to pharmaceutical compositions and/or pharmaceutical preparations comprising at least one compound of the formula I and/or one of its physiologically acceptable salts and/or solvates and especially to pharmaceutical compositions and/or pharmaceutical preparations comprising at least one compound of the formula II and/or one of its physiologically acceptable salts and/or solvates.

- 5
- 10 Preferably, the pharmaceutical compositions and/or pharmaceutical preparations according to the invention contain a therapeutic effective amount of one or more compounds according to the invention. Said therapeutic effective amount of one or more of the compounds according to the invention is known to the skilled artisan or can be easily determined by standard methods known in the art. For example, the compounds according to the invention can be administered to a patient in an analogous manner to other compounds that are effective as raf-kinase inhibitors, especially in an analogous manner to the compounds described in WO 00/42012 (Bayer). Usually, suitable doses that are therapeutically effective lie in the range between 0.0005 mg and 1000 mg, preferably between 0.005 mg and 500 mg and especially between 0.5 and 100 mg per dose unit. The daily dose comprises preferably more than 0.001 mg, more preferred more than 0.01 milligram, even more preferred more than 0.1 mg and especially more than 1.0 mg, for example more than 2.0 mg, more than 5 mg, more than 10 mg, more than 20 mg, more than 50 mg or more than 100 mg, and preferably less than 1500 mg, more preferred less than 750 mg, even more preferred less than 500 mg, for example less than 400 mg, less than 250 mg, less than 150 mg, less than 100 mg, less than 50 mg or less than 10 mg.
- 15
- 20
- 25
- 30

The specific dose for the individual patient depends, however, on the multitude of factors, for example on the efficacy of the specific compounds

- 280 -

employed, on the age, body weight, general state of health, the sex, the kind of diet, on the time and route of administration, on the excretion rate, the kind of administration and the dosage form to be administered, the pharmaceutical combination and severity of the particular disorder to which the therapy relates. The specific therapeutic effective dose for the individual patient can readily be determined by routine experimentation, for example by the doctor or physician which advises or attends the therapeutic treatment.

However, the specific dose for each patient depends on a wide variety of factors, for example on the efficacy of the specific compound employed, on the age, body weight, general state of health, sex, on the diet, on the time and method of administration, on the rate of excretion, medication combination and severity of the particular illness to which the therapy applies. Parenteral administration is preferred. Oral administration is especially preferred.

These compositions and/or preparations can be used as medicaments in human or veterinary medicine. Suitable excipients are organic or inorganic substances which are suitable for enteral (for example oral), parenteral or topical administration and do not react with the novel compounds, for example water, vegetable oils, benzyl alcohols, alkylene glycols, polyethylene glycols, glycerol triacetate, gelatine, carbohydrates, such as lactose or starch, magnesium stearate, talc or vaseline. Examples for suitable dosage forms, which are especially suitable for oral administration are, in particular, tablets, pills, coated tablets, capsules, powders, granules, syrups, juices or drops. Further examples for suitable dosage forms, which are especially suitable for rectal administration are suppositories, further examples for suitable dosage forms, which are especially suitable for parenteral administration are solutions, preferably oil-based or aqueous solutions, furthermore suspensions, emulsions or implants, and suitable for topical application are ointments, creams or

- 281 -

powders. The novel compounds may also be lyophilised and the resultant lyophilisates used, for example, for the preparation of injection preparations. The compositions and/or preparations indicated may be sterilized and/or comprise assistants, such as lubricants, preservatives, stabilizers and/or wetting agents, emulsifiers, salts for modifying the osmotic pressure, buffer substances, dyes and flavors and/or one or more further active ingredients, for example one or more vitamins.

For administration as an inhalation spray, it is possible to use sprays in which the active ingredient is either dissolved or suspended in a propellant gas or propellant gas mixture (for example CO₂ or chlorofluorocarbons). The active ingredient is advantageously used here in micronized form, in which case one or more additional physiologically acceptable solvents may be present, for example ethanol. Inhalation solutions can be administered with the aid of conventional inhalers.

The compounds of the formula I and their physiologically acceptable salts and solvates and especially the compounds of formula II and their physiologically acceptable salts and solvates can be employed for combating one or more diseases, for example allergic diseases, psoriasis and other skin diseases, especially melanoma, autoimmune diseases, such as, for example, rheumatoid arthritis, multiple sclerosis, Crohn's disease, diabetes mellitus or ulcerative colitis.

In General, the substances according to the invention are preferably administered in doses corresponding to the compound rolipram of between 1 and 500 mg, in particular between 5 and 100 mg per dosage unit. The daily dose is preferably between about 0.02 and 10 mg/kg of body weight. However, the specific dose for each patient depends on a wide variety of factors, for example on the efficacy of the specific compound employed, on the age, body weight, general state of health, sex, on the diet, on the time and method of administration, on the excretion rate, medication

combination and severity of the particular illness to which the therapy applies. Oral administration is preferred.

5 The compounds of the formula I according to claim 1 and/or their physiologically acceptable salts are also used in pathological processes which are maintained or propagated by angiogenesis, in particular in tumors, restenoses, diabetic retinopathy, macular degenerative disease or rheumatoid arthritis.

10 Those of skill will readily appreciate that dose levels can vary as a function of the specific compound, the severity of the symptoms and the susceptibility of the subject to side effects. Some of the specific compounds are more potent than others. Preferred dosages for a given compound are readily determinable by those of skill in the art by a variety of means. A preferred means is to measure the physiological potency of a given compound.

15 For use in the subject methods, the subject compounds may be formulated with pharmaceutically active agents other than the compounds according to the invention, particularly other anti-metastatic, antitumor or anti-angiogenic agents. Angiostatic compounds of interest include angiostatin, endostatin, carboxy terminal peptides of collagen alpha (XV), etc. Cytotoxic and cytostatic agents of interest include adriamycin, aleran, Ara-C, BICNU, busulfan, CNUU, cisplatin, cytoxin, daunorubicin, DTIC, 5-FU, hydra, ifosfamide, methotrexate, mithramycin, mitomycin, 20 mitoxantrone, nitrogen mustard, velban, vincristine, vinblastine, VP-16, carboplatinum, fludarabine, gemcitabine, idarubicin, irinotecan, leustatin, navelbine, taxol, taxotere, topotecan, etc.

30 The compounds of the invention have been shown to have antiproliferative effect in an *in vivo* xenograft tumor model. The subject compounds are administered to a subject having a hyperproliferative disorders, e.g., to

5 inhibit tumor growth, to decrease inflammation associated with a lymphoproliferative disorder, to inhibit graft rejection, or neurological damage due to tissue repair, etc. The present compounds are useful for prophylactic or therapeutic purposes. As used herein, the term "treating" is used to refer to both prevention of disease, and treatment of pre-existing conditions. The prevention of proliferation is accomplished by administration of the subject compounds prior to development of overt disease, e.g., to prevent the regrowth of tumors, prevent metastatic growth, diminish restenosis associated with cardiovascular surgery, etc. Alternatively the compounds are used to treat ongoing disease, by stabilizing or improving the clinical symptoms of the patient.

10 The host, or patient, may be from any mammalian species, e.g., primate sp., particularly human; rodents, including mice, rats and hamsters; rabbits; equines, bovines, canines, felines; etc. Animal models are of interest for experimental investigations, providing a model for treatment of human disease.

15 The susceptibility of a particular cell to treatment with the subject compounds may be determined by *in vitro* testing. Typically a culture of the cell is combined with a subject compound at varying concentrations for a period of time sufficient to allow the active agents to induce cell death or inhibit migration, usually between about one hour and one week. For *in vitro* testing, cultured cells from a biopsy sample may be used. The viable cells left after treatment are then counted.

20 The dose will vary depending on the specific compound utilized, specific disorder, patient status, etc. Typically a therapeutic dose will be sufficient to substantially decrease the undesirable cell population in the targeted tissue, while maintaining patient viability. Treatment will generally be continued until there is a substantial reduction, e.g., at least about 50 %, 25 30

- 284 -

decrease in the cell burden, and may be continued until there are essentially none of the undesirable cells detected in the body.

5 The compounds according to the invention are preferably administered to human or nonhuman animals, more preferred to mammalian animals and especially to humans.

10 The compounds also find use in the specific inhibition of a signaling pathway mediated by protein kinases. Protein kinases are involved in signaling pathways for such important cellular activities as responses to extracellular signals and cell cycle checkpoints. Inhibition of specific protein kinases provided a means of intervening in these signaling pathways, for example to block the effect of an extracellular signal, to release a cell from cell cycle checkpoint, etc. Defects in the activity of protein kinases are associated with a variety of pathological or clinical conditions, where there is a defect in the signaling mediated by protein kinases. Such conditions include those associated with defects in cell cycle regulation or in response to extracellular signals, e.g., immunological disorders, autoimmune and immunodeficiency diseases; hyperproliferative disorders, which may include psoriasis, arthritis, inflammation, endometriosis, scarring, cancer, etc. The compounds of the present invention are active in inhibiting purified kinase proteins preferably raf kinases, e.g., there is a decrease in the phosphorylation of a specific substrate in the presence of the compound. The compounds of the invention may also be useful as reagents for studying signal transduction or any of the clinical disorders listed throughout this application.

25 There are many disorders associated with a dysregulation of cellular proliferation. The conditions of interest include, but are not limited to, the following conditions. The subject compounds are useful in the treatment of a variety of conditions where there is proliferation and/or migration of a smooth muscle cells, and/or inflammatory cells into the intimal layer of a

- 285 -

5 vessel, resulting in restricted blood flow through that vessel, e.g., neointimal occlusive lesions. Occlusive vascular conditions of interest include atherosclerosis, graft coronary vascular disease after transplantation, vein graft stenosis, perianastomotic prothetic graft stenosis, restenosis after angioplasty or stent placement, and the like.

10 Diseases where there is hyperproliferation and tissue remodelling or repair or reproductive tissue, e.g., uterine, testicular and ovarian carcinomas, endometriosis, squamous and glandular epithelial carcinomas of the cervix, etc. are reduced in cell number by administration of the subject compounds. The growth and proliferation of neural cells is also of interest.

15 Tumor cells are characterized by uncontrolled growth, invasion to surrounding tissues, and metastatic spread to distant sites. Growth and expansion requires an ability not only to proliferate, but also to down-modulate cell death (apoptosis) and activate angiogenesis to produce a tumor neovasculature.

20 Tumors of interest for treatment include carcinomas, e.g., colon, duodenal, prostate, breast, melanoma, ductal, hepatic, pancreatic, renal, endometrial, stomach, dysplastic oral mucosa, polyposis, invasive oral cancer, non-small cell lung carcinoma, transitional and squamous cell urinary carcinoma etc.; neurological malignancies; e.g. neuroblastoma, gliomas, etc.; hematological malignancies, e.g., childhood acute leukaemia, non-Hodgkin's lymphomas, chronic lymphocytic leukaemia, malignant cutaneous T-cells, mycosis fungoides, non-MF cutaneous T-cell lymphoma, lymphomatoid papulosis, T-cell rich cutaneous lymphoid hyperplasia, bullous pemphigoid, discoid lupus erythematosus, lichen planus, etc.; and the like.

30 Tumors of neural tissue are of particular interest, e.g., gliomas, neuromas, etc. Some cancers of particular interest include breast cancers, which are

primarily adenocarcinoma subtypes. Ductal carcinoma in situ is the most common type of noninvasive breast cancer. In DCIS, the malignant cells have not metastasized through the walls of the ducts into the fatty tissue of the breast. Infiltration (or invasive) ductal carcinoma (IDC) has metastasized through the wall of the duct and invaded the fatty tissue of the breast. Infiltrating (or invasive) lobular carcinoma (ILC) is similar to IDC, in that it has the potential to metastasize elsewhere in the body. About 10 % to 15 % of invasive breast cancers are invasive lobular carcinomas.

5

10

Also of interest is non-small cell lung carcinoma. Non-small cell lung cancer (NSCLC) is made up of three general subtypes of lung cancer. Epidermoid carcinoma (also called squamous cell carcinoma) usually starts in one of the larger bronchial tubes and grows relatively slowly. The size of these tumors can range from very small to quite large. Adenocarcinoma starts growing near the outside surface of the lung and may vary in both size and growth rate. Some slowly growing adenocarcinomas are described as alveolar cell cancer. Large cell carcinoma starts near the surface of the lung, grows rapidly, and the growth is usually fairly large when diagnosed. Other less common forms of lung cancer are carcinosarcoma, mucoepidermoid, and malignant mesothelioma.

15

20

Melanoma is a malignant tumor of melanocytes. Although most melanomas arise in the skin, they also may arise from mucosal surfaces or at other sites to which neural crest cells migrate. Melanoma occurs predominantly in adults, and more than half of the cases arise in apparently normal areas of the skin. Prognosis is affected by clinical and histological factors and by anatomic location of the lesion. Thickness and/or level of invasion of the melanoma, mitotic index, tumor infiltrating lymphocytes, and ulceration or bleeding at the primary site affect the prognosis. Clinical staging is based on whether the tumor has spread to regional lymph nodes or distant sites. For disease clinically confined to the

30

primary site, the greater the thickness and depth of local invasion of the melanoma, the higher the chance of lymph node metastases and the worse the prognosis. Melanoma can spread by local extension (through lymphatics) and/or by hematogenous routes to distant sites. Any organ may be involved by metastases, but lungs and liver are common sites.

5

Other hyperproliferative diseases of interest relate to epidermal hyperproliferation, tissue, remodeling and repair. For example, the chronic skin inflammation of psoriasis is associated with hyperplastic epidermal keratinocytes as well as infiltrating mononuclear cells, including CD4+ memory T cells, neutrophils and macrophages.

10

The proliferation of immune cells is associated with a number of autoimmune and lymphoproliferative disorders. Diseases of interest include multiple sclerosis, rheumatoid arthritis and insulin dependent diabetes mellitus. Evidence suggests that abnormalities in apoptosis play a part in the pathogenesis of systemic lupus erythematosus (SLE). Other lymphoproliferative conditions the inherited disorder of lymphocyte apoptosis, which is an autoimmune lymphoproliferative syndrome, as well as a number of leukemia's and lymphomas. Symptoms of allergies to environmental and food agents, as well as inflammatory bowel disease, may also be alleviated by the compounds of the invention.

15

20

Surprisingly, it has been found that methylene urea derivatives according to invention are able to interact with signaling pathways, especially the signaling pathways described herein and preferably the raf-kinase signaling pathway. Methylene urea derivatives according to the invention preferably show advantageous biological activity which can easily be demonstrated according to methods known in the art, for example by enzyme based assays. Suitable assays are known in the art, for example from the literature cited herein and the references cited in the literature, or can be developed and/or performed in an analogous manner thereof. In

25

30

- 288 -

5 such enzyme based assays, methylene urea derivatives according to invention show an effect, preferably a modulating and especially an inhibiting effect which is usually documented by IC_{50} values in a suitable range, preferably in the micromolar range and more preferred in the nanomolar range.

10 In general, compounds according to the invention are to be regarded as suitable kinase-modulators and especially suitable kinase-inhibitors according to the invention if they show an effect or an activity to one or more kinases, preferably to one or more raf-kinases that preferably lies, determined as IC_{50} -value, in the range of 100 μ mol or below, preferably 10 μ mol or below, more preferably in the range of 3 μ mol or below, even more preferably in the range of 1 μ mol or below and most preferably in the nanomolar range. Especially preferred for use according to the invention are kinase-inhibitors as defined above/below, that show an activity, determined as IC_{50} -value, to one or more raf-kinases, preferably including A-raf, B-raf and c-raf1 or consisting of A-raf, B-raf and c-raf1 and more preferred including c-raf1 or consisting of c-raf1, in the range of 0.5 μ mol or below and especially in the range of 0.1 μ mol or below. In many cases an IC_{50} -value at the lower end of the given ranges is advantageous and in some cases it is highly desirable that the IC_{50} -value is as small as possible or the IC_{50} -values are as small as possible, but in general IC_{50} -values that lie between the above given upper limits and a lower limit in the range of 0.0001 μ mol, 0.001 μ mol, 0.01 μ mol or even above 0.1 μ mol are sufficient to indicate the desired pharmaceutical activity. However, the activities measured can vary depending on the respective testing system or assay chosen.

15 Alternatively, the advantageous biological activity of the compounds according to the invention can easily be demonstrated in *in vitro* assays, such as *in vitro* proliferation assays or *in vitro* growth assays. Suitable *in vitro* assays are known in the art, for example from the literature cited

- 289 -

5 herein and the references cited in the literature or can be performed as described below, or can be developed and/or performed in an analogous manner thereof.

10 As an example for an *in vitro* growth assay, human tumor cell lines, for example HCT116, DLD-1 or MlaPaCa, containing mutated K-ras genes can be used in standard proliferation assays, for example for anchorage dependent growth on plastic or anchorage independent growth in soft agar. Human tumor cell lines are commercially available, for example from ATCC (Rockville MD), and can be cultured according to methods known in the art, for example in RPMI with 10% heat inactivated fetal bovine serum and 200 mM glutamine. Cell culture media, fetal bovine serum and additives are commercially available, for example from Invitrogen/Gibco/BRL (Karlsruhe, Germany) and/or QRH Biosciences (Lenexa, KS). In a standard proliferation assay for anchorage dependent growth, 3×10^3 cells can be seeded into 96-well tissue culture plates and allowed to attach, for example overnight at 37 °C in a 5% CO₂ incubator. Compounds can be titrated in media in dilution series and added to 96 well cell cultures. Cells are allowed to grow, for example for 1 to 5 days, typically with a feeding of fresh compound containing media at about half of the time of the growing period, for example on day 3, if the cells are allowed to grow 5 days. Proliferation can be monitored by methods known in the art, such as measuring metabolic activity, for example with standard XTT colorimetric assay (Boehringer Mannheim) measured by standard ELISA plate reader at OD 490/560, by measuring 3 H-thymidine incorporation into DNA following an 8 h culture with 1 μ Ci 3 H-thymidine, harvesting the cells onto glass fiber mats using a cell harvester and measuring 3 H-thymidine incorporation by liquid scintillation counting, or by staining techniques, such as crystal violet staining. Other suitable cellular assay systems are known in the art.

- 290 -

Alternatively, for anchorage independent cell growth, cells can be plated at 1×10^3 to 3×10^3 in 0.4% Seaplaque agarose in RPMI complete media, overlaying a bottom layer containing only 0.64% agar in RPMI complete media, for example in 24-well tissue culture plates. Complete media plus dilution series of compounds can be added to wells and incubated, for example at 37 °C in a 5% CO₂ incubator for a sufficient time, for example 10-14 days, preferably with repeated feedings of fresh media containing compound, typically at 3-4 day intervals. Colony formation and total cell mass can be monitored, average colony size and number of colonies can be quantitated according to methods known in the art, for example using image capture technology and image analysis software. Image capture technology and image analysis software, such as Image Pro Plus or media Cybertetics.

As discussed herein, these signaling pathways are relevant for various disorders. Accordingly, by interacting with one or more of said signaling pathways, methylene urea derivatives are useful in the prevention and/or the treatment of disorders that are dependent from said signaling pathways.

The compounds according to the invention are preferably kinase modulators and more preferably kinase inhibitors. According to the invention, kinases include, but are not limited to one or more Raf-kinases, one or more Tie-kinases, one or more VEGFR-kinases, one or more PDGFR-kinases, p38-kinase and/or SAPK2alpha.

Raf-kinases in this respect are respect preferably include or consist of A-Raf, B-Raf and c-Raf1.

Tie-kinases in this respect preferably include or consist of Tie-2 kinase.

- 291 -

VEGFR-kinases in this respect preferably include or consist of VEGFR-2 kinase.

Due to the kinase modulating or inhibiting properties of the compounds according to the invention, the compounds according to the invention preferably interact with one or more signalling pathways which are preferably cell signalling pathways, preferably by downregulating or inhibiting said signalling pathways. Examples for such signalling pathways include, but are not limited to the raf-kinase pathway, the Tie-kinase pathway, the VEGFR-kinase pathway, the PDGFR-kinase pathway, the p38-kinase pathway, the SAPK2alpha pathway and/or the Ras-pathway.

Modulation of the raf-kinase pathway plays an important role in various cancerous and noncancerous disorders, preferably cancerous disorders, such as dermatological tumors, haematological tumors, sarcomas, squamous cell cancer, gastric cancer, head cancer, neck cancer, oesophageal cancer, lymphoma, ovary cancer, uterine cancer and/or prostate cancer. Modulation of the raf-kinase pathway plays a even more important role in various cancer types which show a constitutive activation of the raf-kinase dependent signalling pathway, such as melanoma, colorectal cancer, lung cancer, brain cancer, pancreatic cancer, breast cancer, gynaecological cancer, ovarian cancer, thyroid cancer, chronic leukaemia and acute leukaemia, bladder cancer, hepatic cancer and/or renal cancer. Modulation of the raf-kinase pathway plays also an important role in infection diseases, preferably the infection diseases as mentioned above/below and especially in Helicobacter pylori infections, such as Helicobacter pylori infection during peptic ulcer disease.

One or more of the signalling pathways mentioned above/below and especially the VEGFR-kinase pathway plays an important role in angiogenesis. Accordingly, due to the kinase modulating or inhibiting properties of the compounds according to the invention, the compounds

- 292 -

5 according to the invention are suitable for the prophylaxis and/or treatment of pathological processes or disorders caused, mediated and/or propagated by angiogenesis, for example by inducing anti-angiogenesis. Pathological processes or disorders caused, mediated and/or propagated by angiogenesis include, but are not limited to tumors, especially solid tumors, arthritis, especially rheumatoid arthritis, diabetic retinopathy, psoriasis, restenosis, fibrotic disorders, mesangial cell proliferative disorders, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, organ transplant rejection, glomerulopathies, metabolic disorders, inflammation and

10 neurodegenerative diseases, and especially solid tumors, rheumatic arthritis, diabetic retinopathy and psoriasis.

15 Modulation of the p38-signalling pathway plays an important role in various cancerous and although in various noncancerous disorders, such as fibrosis, atherosclerosis, restenosis, vascular disease, cardiovascular disease, inflammation, renal disease and/or angiogenesis, and especially noncancerous disorders such as rheumatoid arthritis, inflammation, autoimmune disease, chronic obstructive pulmonary disease, asthma and/or inflammatory bowel disease.

20

Modulation of the PDGF-signalling pathway plays an important role in various cancerous and although in various noncancerous disorders, such as rheumatoid arthritis, inflammation, autoimmune disease, chronic obstructive pulmonary disease, asthma and/or inflammatory bowel disease, and especially noncancerous disorders such as fibrosis, atherosclerosis, restenosis, vascular disease, cardiovascular disease, inflammation, renal disease and/or angiogenesis.

25

30 Subject of the present invention are therefore methylene urea derivatives according to the invention as promoters or inhibitors, preferably as inhibitors, of the signalling pathways described herein. Preferred subject of

- 293 -

5 the invention are therefore methylene urea derivatives according to the invention as promoters or inhibitors, preferably as inhibitors of the raf-kinase pathway. More preferred subject of the invention are therefore methylene urea derivatives according to the invention as promoters or inhibitors, preferably as inhibitors of the raf-kinase. Even more preferred subject of the invention are methylene urea derivatives according to invention as promoters or inhibitors, preferably as inhibitors of one or more raf-kinases, selected from the group consisting of A-raf, B-raf and c-raf1. Especially preferred subject of the invention are methylene urea derivatives according to the invention as promoters or inhibitors, preferably as inhibitors of c-raf1.

10

15 Thus, subject of the present invention are methylene urea derivatives according to the invention as medicaments. Subject of the present invention are methylene urea derivatives according to the invention as medicament active ingredients. Further subject of the present invention is the use of one or more methylene urea derivatives according to the invention as a pharmaceutical. Further subject of the present invention is the use of one or more methylene urea derivatives according to the invention in the treatment and/or the prophylaxis of disorders, preferably the disorders described herein, more preferred disorders that are caused, mediated and/or propagated by signalling pathways discussed herein, even more preferred disorders that are caused, mediated and/or propagated by raf-kinases and especially disorders that are caused, mediated and/or propagated by raf-kinases, selected from the group consisting of A-raf, B-raf and c-raf1. Usually, the disorders discussed herein are divided into two groups, hyperproliferative and non hyperproliferative disorders. In this context, psoriasis, arthritis, inflammation, endometriosis, scarring, benign prostatic hyperplasia, immunological diseases, autoimmune diseases and immunodeficiency diseases are to be regarded as noncancerous disorders, of which arthritis, inflammation, immunological diseases, autoimmune diseases and

25

30

5 immunodeficiency diseases are usually regarded as non hyperproliferative disorders. In this context, brain cancer, lung cancer, squamous cell cancer, bladder cancer, gastric cancer, pancreatic cancer, hepatic cancer, renal cancer, colorectal cancer, breast cancer, head cancer, neck cancer, oesophageal cancer, gynaecological cancer, thyroid cancer, lymphoma, chronic leukaemia and acute leukaemia are to be regarded as cancerous disorders, all of which are usually regarded as hyperproliferative disorders. Especially cancerous cell growth and especially cancerous cell growth mediated by raf-kinase is a disorder which is a target of the present invention. Subject of the present invention therefore are methylene urea derivatives according to the invention as medicaments and/or medicament active ingredients in the treatment and/or the prophylaxis of said disorders and the use of methylene urea derivatives according to the invention for the manufacture of a pharmaceutical for the treatment and/or the prophylaxis of said disorders as well as a method of treatment of said disorders, comprising administering one or more methylene urea derivatives according to the invention to a patient in need of such an administration. Subject of the present invention therefore are methylene urea derivatives according to the invention as medicaments and/or medicament active ingredients in the treatment and/or the prophylaxis said disorders and the use of methylene urea derivatives according to the invention for the manufacture of a pharmaceutical for the treatment and/or the prophylaxis of said disorders as well as a method of treatment of said disorders, comprising administering one or more methylene urea derivatives according to the invention to a patient in need of such an administration.

25

Accordingly, subject of the present invention are pharmaceutical compositions that contain one or more methylene urea derivatives according to the invention. Subject of the present invention are especially pharmaceutical compositions that contain one or more methylene urea derivatives according to the invention and one or more additional

30

compounds (other than the compounds of the instant invention), preferably selected from the group consisting of physiologically acceptable excipients, auxiliaries, adjuvants, carriers and pharmaceutically active ingredients other than the compounds according to the invention.

5 Accordingly, subject of the present invention is a process for the manufacture of a pharmaceutical composition, wherein one or more methylene urea derivatives according to the invention and one or more compounds (other than the compounds of the instant invention), preferably selected from the group consisting of carriers, excipients, auxiliaries, adjuvants and pharmaceutically active ingredients other than the compounds according to the invention.

10

Accordingly, the use of the compounds according to the invention in the treatment of Hyperproliferative disorders is a subject of the instant invention.

15

Accordingly, the use of the compounds according to the invention for producing a medicament for the treatment of hyperproliferative disorders is a subject of the instant invention.

20

Above and below, all temperatures are given in °C. In the examples below, "conventional work-up" means that the organic phase is washed with saturated NaHCO₃ solution, if desired with water and saturated NaCl solution, the phases are separated, the organic phase is dried over sodium sulfate and evaporated, and the product is purified by chromatography on silica gel, by preparative HPLC and/or by crystallization.

25

The present invention relates to methylene urea derivatives of formula I, the use of the compounds of formula I as inhibitors of raf-kinase, the use of the compounds of formula I for the manufacture of a pharmaceutical composition and a method of treatment, comprising administering said pharmaceutical composition to a patient.

30


- 296 -

Examples

Experimental part

5 Synthesis of the benzylamine moieties

4-(4-Pyridinyloxy)benzylamine




- 15 a) 5 g (42 mmol) of 4-hydroxybenzonitrile and 13.36 g (42 mmol) of bipyrindine are mixed and heated to 150°C. After the reaction mixture has been stirred at 150°C for 3 hours, it is cooled, 500 ml of 10% Na₂CO₃ solution are added, and the mixture is stirred. The resultant precipitate is filtered off with suction, rinsed with 500 ml of water and dried under reduced pressure. Extraction of the aqueous phase with ethyl acetate followed by drying and evaporation gave further product.
- 20 Yield: 3.86 g (47%) of 1, pale-brown solid

- 25 b) Compound 1 is hydrogenated using Raney nickel in methanolic ammonia solution at 50°C and 5 bar. The reaction solution is filtered through Kieselguhr and rinsed with MeOH, and the filtrate is subsequently evaporated.
- 30 Yield: 3.49 g (78%) of 2, brown oil

- 297 -


3-(4-Pyridinyloxy)benzylamine



- 10 a) 5 g (42 mmol) of 3-hydroxybenzonitrile and 13.36 g (42 mmol) of bipyrindine are mixed and heated to 150°C. After the reaction mixture has been stirred at 150°C for 3 hours, it is cooled, diluted with 600 ml of ethyl acetate and washed with 600 ml of 10% Na₂CO₃ solution. The organic phase is dried using Na₂SO₄, filtered and evaporated. The residue is purified by column chromatography (100 g of silica gel; eluent: ethyl acetate:petroleum ether = 1:1).
- 15 Yield: 2.46 g (35%) of 3, pale-yellow crystals

- 20 b) Compound 3 is hydrogenated using Raney nickel in methanolic ammonia solution at 50°C and 5 bar. The reaction solution is filtered through Kieselguhr and rinsed with MeOH, and the filtrate is subsequently evaporated.
- 25 Yield: 2.33 g (96%) of 4, brown oil

4-(3-Pyridinyloxy)benzylamine



- a) 3 g (25 mmol) of 4-hydroxybenzonitrile; 2.44 g (25 mmol) of 3-bromopyridine and 5.71 g (30 mmol) of copper iodide are dissolved in DMF, 6.91 g (50 mmol) of K_2CO_3 are added, and the mixture is refluxed for 18 hours. The reaction mixture is cooled, 200 ml of dichloromethane are added, and the mixture is stirred for 15 minutes and filtered. The filtrate is washed with water and extracted with 10% HCl solution. The HCl phase is neutralized using NH_4OH solution and extracted with dichloromethane. The combined organic phases are washed with $NaOH$ solution (2M), dried using Na_2SO_4 , filtered and evaporated. The residue is purified by column chromatography (25 g of silica gel, eluent: ethyl acetate:petroleum ether = 1:2).


5

Yield: 375 mg (8%) of **5**, yellow crystals

10

- b) Compound **5** is hydrogenated using Raney nickel in methanolic ammonia solution at 50°C and 4.8 bar. The reaction solution is filtered through kieselguhr and rinsed with MeOH, and the filtrate is subsequently evaporated.

15

Yield: 440 mg (97%) of **6**, brown oil**3-(3-Pyridinyloxy)benzylamine**

25

- a) 1 g (11 mmol) of 3-hydroxypyridine and 3.26 g (22 mmol) of 3-nitrobenzonitrile are dissolved in DMF, 3.34 g (24 mmol) of potassium carbonate are added, and the mixture is refluxed overnight. The reaction mixture is evaporated, and the residue is taken up in 150 ml of dichloromethane, stirred for 30 minutes and filtered. The filtrate is washed

30

with water and extracted with 10% HCl solution. The HCl phase is neutralized using NH_4OH and extracted with dichloromethane. The combined organic phases are dried using Na_2SO_4 , filtered and evaporated. The residue is purified by column chromatography (33 g of silica gel, eluent: ethyl acetate:petroleum ether = 1:2).

5


Yield: 956 mg (44%) of **7**, yellow oil

- b) Compound **7** is hydrogenated using Raney nickel in methanolic ammonia solution at 50°C and 4.8 bar. The reaction solution is filtered through kieselguhr and rinsed with MeOH, and the filtrate is subsequently evaporated.

10

Yield: 945 mg (97%) of **8**, brown oilSynthesis of the benzylureas

15

Variant A**1-(4-Chloro-3-trifluoromethylphenyl)-3-{4-(4-pyridinyloxy)benzyl}urea**

20


150 mg (0.75 mmol) of **2** are dissolved in dichloromethane together with 166 mg (0.75 mmol) of 4-chloro-3-trifluoromethylphenyl isocyanate, and the mixture is stirred at room temperature for 2 hours. The reaction mixture is evaporated, and the residue is purified by column chromatography (11 g of silica gel, eluent: ethyl acetate:petroleum ether = 2:1).

25


Yield: 119 mg (37%), colourless oil

30

- 300 -


1-(4-Chloro-3-trifluoromethylphenyl)-3-[3-(4-pyridinyloxy)benzyl]urea

100 mg (0.5 mmol) of **4** are dissolved in dichloromethane together with 133 mg (0.6 mmol) of 4-chloro-3-trifluoromethylphenyl isocyanate and 0.1 ml (0.6 mmol) of N-ethyl-diisopropylamine, and the mixture is stirred at room temperature for 2 hours. The resultant precipitate is filtered off with suction, washed with dichloromethane and dried under reduced pressure. Yield: 201 mg (96%), colourless solid


1-(4-Chloro-3-trifluoromethylphenyl)-3-[4-(3-pyridinyloxy)benzyl]urea

100 mg (0.5 mmol) of **6** are dissolved in dichloromethane together with 111 mg (0.5 mmol) of 4-chloro-3-trifluoromethylphenyl isocyanate. After the reaction mixture has been stirred at room temperature for 4 hours, it is refluxed for 1 hour and subsequently stirred again at room temperature overnight. The reaction mixture is diluted with dichloromethane, extracted with saturated Na_2HCO_3 solution, dried using Na_2SO_4 , filtered and evaporated. The residue is purified by preparative HPLC. Yield: 40 mg (19%), yellow oil

- 301 -

1-(4-Chloro-3-trifluoromethylphenyl)-3-[3-(3-pyridinyloxy)benzyl]urea


100 mg (0.5 mmol) of **8** are dissolved in dichloromethane together with 111 mg (0.5 mmol) of 4-chloro-3-trifluoromethylphenyl isocyanate. After the reaction mixture has been stirred at room temperature for 4 hours, it is refluxed for 1 hour and subsequently stirred again at room temperature overnight. The reaction mixture is diluted with dichloromethane, extracted with saturated Na_2HCO_3 solution, dried using Na_2SO_4 , filtered and evaporated. The residue is purified by column chromatography (3 g of silica gel, eluent: ethyl acetate:petroleum ether = 1:3). Yield: 34 mg (16%), colourless solid

Variant B**5-(4-Butyl-3-isoxazolyl)carbamic acid, 4-nitrophenyl ester**

2.91 g (20.76 mmol) of 3-amino-5-tert-butylisoxazole and 1.84 ml (22.83 mmol) of pyridine are dissolved in dichloromethane, and 4.18 g (20.76 mmol) of 4-nitrophenyl chloroformate are added at room temperature. After the reaction mixture has been stirred at room temperature for 2.5 hours, it is evaporated, and the residue is digested with diethyl ether, filtered off with suction and subsequently dried under reduced pressure. Yield: 5.68 g (90%) of **9**, colourless solid

- 302 -

1-(5-t-Butyl-3-isoxazolyl)-3-[4-(4-pyridinyloxy)benzyl]urea




100 mg (0.33 mmol) of **9** and 79 mg (0.39 mmol) of **2** are dissolved in dichloromethane, and the solution is stirred at room temperature for 4 hours. The reaction mixture is evaporated, and the residue is purified by column chromatography (5 g of silica gel, eluent: ethyl acetate).

Yield: 67 mg (56%), colourless solid

10

1-(5-t-Butyl-3-isoxazolyl)-3-[3-(4-pyridinyloxy)benzyl]urea



100 mg (0.33 mmol) of **9** and 79 mg (0.39 mmol) of **4** are dissolved in dichloromethane, and the solution is stirred at room temperature for 4 hours. The reaction mixture is evaporated, and the residue is purified by column chromatography (5 g of silica gel, eluent: ethyl acetate).

Yield: 33 mg (27.5%), colourless oil


20

Synthesis of the substituted benzyl amino building blocks

25

- 303 -

4-[(4-Aminomethyl)phenoxy]-2-pyridine carboxylic acid, methylamide



10

15

60 ml Thionylchloride are heated to a temperature of 45 °C under a nitrogen atmosphere and 1.83 ml Dimethylformamide is added slowly. 20 g Pyridin-2-carboxylic acid is added to the solution in portions, the reaction mixture is stirred another 15 min at 45 °C and then heated to 80 °C for 24 hrs. The reaction mixture is evaporated and the resulting residue treated with dry toluene as a carrier and then evaporated. This procedure is repeated several times. The resulting oil is dissolved in toluene, cooled to 0 °C, slowly treated with methanol and stirred for one hour. The resulting precipitate is filtered by suction, washed with toluene and recrystallised from acetone.

25

Yield: 15 g (44 %) **10**, colourless crystals

30

13 g (62.5 mmol) **10** are dissolved together with 2.98 g (31.24 mmol) dry Magnesiumchloride in THF. After 5 min 110 ml methyl amine solution (2M

- 304 -

in THF) are added dropwise within 10 min and the suspension stirred for 2 h at room temperature. 120 ml water und 63 ml 1M HCl- solution are added and the mixture is extracted three times with ethyl acetate. The combined organic phases are washed with brine, dried with Na_2SO_4 , filtered and evaporated.

Yield: 10.5 g (98.5 %) 11, colourless oil.


4.15 g (24.32 mmol) 11 are heated to a temperature of 160 °C together with 5.8 g (48.65 mmol) 4-Cyanophenol for 18 hrs in an argon atmosphere. The reaction mixture is cooled down, diluted with ethyl acetate, washed consecutively twice with 30 ml 2N NaOH- solution, twice with 30 ml Wasser and once with 30 ml brine, dried over Na_2SO_4 , filtered and evaporated. The residue is digested with diethyl ether, filtered by suction, washed with diethyl ether:petrol ether = 1:1 and dried in vacuo.

Yield: 3.27 g (52 %) 12, brownish solid.

3.27 g (12.65 mmol) 12 are hydrogenated in methanolic ammonia solution in the presence of Raney nickel at 45 °C and 5 bar pressure. The reaction mixture is filtered over kieselguhr, washed with MeOH and the filtrate is evaporated. The residue is purified by chromatography (120 g kieselgel, eluent: CH_2Cl_2 / methanol/ NH_3 (9:1 + 0, 1%).

Yield: 2.55 g (78 %) 13, yellow solid

4-[(3-Aminomethyl)phenoxy]-2-pyridincarbonsäure, methylamid



- 305 -

5 g (29.31 mmol) 11 are heated together with 6.98 g (58.62 mmol) 3-Cyanophenol at a temperature of 160 °C in an argon atmosphere. After 18 hrs, further 3.49 g (29.30 mmol) 3-Cyanophenol are added and heating is continued for 6 hrs at 160 °C. The reaction mixture is cooled, diluted with ethyl acetate, washed consecutively twice with 40 ml 2N NaOH- solution, twice with 35 ml water and once with 30 ml brine, dried over Na_2SO_4 , filtered and evaporated. The residue is digested with diethyl ether, filtered by suction, washed with diethyl ether:petrol ether = 2:1 and dried in vacuo. Another portion of product is obtained by chromatography of the mother liquor (95 g silica gel, eluent: ethyl acetat:petrol ether = 7:3).


Yield: 3.58 g (46 %) 14, brownish solid

3.16 g (12.65 mmol) 14 are hydrogenated in methanolic ammonia solution in the presence of Raney nickel at 45 °C and 5 bar pressure. The reaction mixture is filtered over kieselguhr, washed with MeOH and the filtrate is evaporated. The residue is purified by chromatography (120 g silica gel, eluent: CH_2Cl_2 / methanol/ NH_3 (9:1 + 0, 1%).

Yield: 2.67 g (86 %) 15, light brownish oil

Synthesis of benzyl ureas

Variant A



A solution of 0.16 mmol isocyanate and 0.16 mmol benzyl amine 2 or 13 in dichloromethane is stirred 16 h at room temperature. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

- 306 -

Variant A: The resulting precipitate is filtered by suction, washed consecutively with dichloromethane, ethyl acetate and diethyl ether and dried in vacuo at 40 °C.


Variant B: The reaction mixture is evaporated and the residue is treated with 0.5 ml acetonitrile. The resulting precipitate is filtered by suction, washed with acetonitrile and diethyl ether and dried in vacuo at 40 °C.

Variant C: The reaction mixture is evaporated, the oily residue taken up in 2 ml acetonitrile: water = 1:1, frozen and then freeze dried for 16 hrs.

Variant D: The reaction mixture is evaporated to dryness. The residue is dried in vacuo at 40 °C.

Variant E: The reaction mixture is evaporated to dryness. The residue is purified by chromatography (4 g silica gel, eluent: dichloromethane:methanol = 98:2 to 95:5). The obtained crude product is taken up in 1.2 ml acetonitrile:water = 2:1, frozen and freeze dried for 16 hrs.

Variant F: The reaction mixture is evaporated to dryness, the residue dissolved in 15 ml ethyl acetate and extracted once with 10 ml 25 % hydrochloric acid. The water phase is separated, made alkaline with 32 % sodium hydroxide solution (pH= 9-10) and extracted three times, each time with 10 ml ethyl acetate. The combined organic phases are washed with 15 ml brine, dried over Na₂SO₄, filtered and the filtrate evaporated. The residue is dried at 40 °C overnight.



0.16 mmol isocyanate and 0.16 mmol benzyl amine **4** or **15** are dissolved in dichloromethane gelöst and stirred for 16 hrs at room temperature. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

- 307 -

Variant A: The resulting precipitate is filtered by suction, washed consecutively with dichloromethane, ethyl acetate and diethyl ether and dried in vacuo at 40 °C.

Variant B: The reaction mixture is evaporated. The residue is dried in vacuo at 40 °C.

Variant C: The reaction mixture is evaporated to dryness. The residue is purified by chromatography (4 g silica gel, eluent: dichloromethane:methanol = 98:2 bis 95:5). The obtained crude product is taken up in 1.2 ml acetonitrile:water = 2:1, frozen and freeze dried for 16 hrs.


Variant D: The reaction mixture is evaporated, the oily residue taken up in 2 ml acetonitrile:water = 1:1, frozen and freeze dried for 16 hrs.

Variant E: The reaction mixture is evaporated, the residue digested with 1 ml ethyl acetate:diethyl ether = 2:1, filtered by suction, washed with diethyl ether and dried in vacuo at 40 °C.

Variant F: The reaction mixture is evaporated and the residue is treated with 0.5 ml acetonitrile. The resulting precipitate is filtered by suction, washed with acetonitrile and diethyl ether and dried in vacuo at 40 °C.

Variant G: The reaction mixture is evaporated to dryness, the residue dissolved in 15 ml ethyl acetate and extracted once with 10 ml 25 % hydrochloric acid. The water phase is separated, made alkaline with 32 % sodium hydroxide solution (pH= 9-10) and extracted three times, each time with 10 ml ethyl acetate. The combined organic phases are washed with 15 ml brine, dried over Na₂SO₄, filtered and the filtrate evaporated. The residue is dried at 40 °C overnight.


- 308 -

Variant Bb) synthesis of the trichloro aceto anilides

10 1 g of the substituted aniline and 1.3 eq. pyridine are dissolved in dichloromethane, cooled to 0 °C and 1.1 eq. trichloro acetic acid chloride is slowly added. After the addition is completed, the reaction mixture is allowed warm up to room temperature and stirring is continued for 1 h. Then the reaction mixture is extracted 1N hydrochloric acid and water consecutively and the organic phase is dried over Na₂SO₄, filtered and evaporated.

- 15
- R₁ = 2-OMe, 5-CF₃; colourless solid, yield: 93 %
 R₁ = 3-CF₃, 4-Br; colourless solid, yield: 100 %
 R₁ = 3-OCF₃; yellow solid, yield: 82 %
 R₁ = 2-OMe, 4-Me, 5-Cl; beige solid, yield: 84 %
 R₁ = 2-OMe, 4-Cl, 5-CF₃; yellow oil, either: 85 %
 R₁ = 2-S-Me, 5-CF₃; yellow solid, yield: 92 %
 R₁ = 2-OMe, 5-Me; beige solid, yield: 99 %
 R₁ = 3-Me, 4-Cl; colourless solid, yield: 97 %

25 b) synthesis of the final products



- 309 -


0.15 mmol of the substituted trichloro aceto anilide and 0.15 mmol benzyl amine **2** or **13** are dissolved in DMSO, 0.15 mmol DBU are added and the mixture heated to 80 °C for 2.5 - 5.5 hrs. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

5 Variant A: The reaction mixture is cooled, diluted with dichloromethane and extracted with 1N hydrochloric acid. The water phase is made alkaline with 2N sodium hydroxide solution and extracted several times with ethyl acetate. The combined organic phases are dried over Na₂SO₄, filtered and evaporated.

10 Variant B: The reaction mixture is cooled, diluted with dichloromethane and extracted with 1N hydrochloric acid. The water phase is made alkaline with 2N sodium hydroxide solution and extracted several times with ethyl acetate. The combined organic phases are dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (5 g silica gel, eluent: dichloromethane:methanol = 98:2 to 95:5).

15 Variant C: The reaction mixture is cooled, diluted with dichloromethane and washed with water. The organic phases what dried over Na₂SO₄, filtered and evaporated. The residue is digested with a small amount of water, the resulting precipitate filtered by suction and dried in vacuo at 40 °C.

20 Variant D: The reaction mixture is cooled, diluted with dichloromethane, extracted consecutively twice with 1N hydrochloric acid and with water, dried over Na₂SO₄, filtered and evaporated.



- 310 -

mixture heated to 80 °C for 2.5 - 5.5 hrs. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

Variant A: The reaction mixture is cooled, diluted with dichloromethane and extracted with 1N hydrochloric acid. The water phase is made alkaline with 2N sodium hydroxide solution and extracted several times with ethyl acetate. The combined organic phases are dried over Na₂SO₄, filtered and evaporated.


Variant B: The reaction mixture is cooled, diluted with dichloromethane and extracted with 1N hydrochloric acid. The water phase is made alkaline with 2N sodium hydroxide solution and extracted several times with ethyl acetate. The combined organic phases are dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (5 g silica gel, eluent:dichloromethane:methanol = 98:2 to 95:5).

Variant C: The reaction mixture is cooled, diluted with dichloromethane and washed with water. The organic phases are dried over Na₂SO₄, filtered and evaporated. The residue is digested with a small amount of water, the resulting precipitate filtered by suction and dried in vacuo at 40 °C.

20

Variant C

a) Synthesis of the anilines



30

- 311 -

2 mmol 4-Fluoro-3-nitrobenzotrifluoride are dissolved in dimethyl sulfoxide (DMSO), treated with 2-2.4 mmol amine and stirred at 50 °C overnight. The reaction mixture is diluted with dichloromethane and extracted twice with water. The organic phase is dried over Na₂SO₄, filtered and evaporated in vacuo. The obtained crude product is employed in the next synthesis step without further purification.

R₁, R₂ = Me; orange oil, yield: 96 %

R₁, R₂ = Et; orange oil, yield: 97 %

R₁ = Me, R₂ = (CH₂)₂OCH₃; orange oil, yield: 91.5 %

R₁ = Me, R₂ = (CH₂)₂N(CH₃)₂; orange oil, yield: 85 %

The accordingly obtained nitro compounds are hydrogenated in THF in the presence of H₂ and Pd/C (5%) at room temperature overnight. Then the catalyst is separated by filtration and the filtrate evaporated to yield the respective aniline.


R₁, R₂ = Me; yellow oil, yield: 92 %

R₁, R₂ = Et; yellow oil, yield: 92 %

R₁ = Me, R₂ = (CH₂)₂OCH₃; red oil, yield: 99 %

R₁ = Me, R₂ = (CH₂)₂N(CH₃)₂; yellow oil, yield 98.5 %

R₁ = Me, R₂ = (CH₂)₂N(CH₃)₂; yellow oil, yield: 98.5 %



30


- 312 -

2 mmol 4-Fluor-3-nitrobenzotrifluoride are dissolved in dimethylformamide, treated with 2.2 mmol 2-(Dimethylamino)ethanol and 4.6 mmol cesium carbonate and stirred at room temperature. After 48 h, additional 1 mmol 2-(Dimethylamino)ethanol are added and at the reaction mixture is stirred at 40 °C overnight. The reaction mixture is diluted with ethyl acetate and the resulting solution washed twice with water. The organic phase is dried over Na₂SO₄, filtered, treated with dry toluene as a carrier and then evaporated several times using a Rotavapor and then evaporated for dryness. The obtained residue is purified by chromatography (35 g silica gel, eluent: dichloromethane:acetone = 100:0 to 90:10).

Yield: 43 %, yellow oil

The accordingly obtained nitro compound is hydrogenated in THF in the presence of H₂ and Pd/C (5%) at room temperature overnight. Then the catalyst is separated by filtration and the filtrate evaporated to yield the respective aniline.

Yield: 97 %, yellow crystals




5-Chlor-2-nitroanisol in THF are hydrogenated in the presence of H₂ and Raney nickel at room temperature overnight. The catalyst is separated by filtration and the filtrate evaporated to dryness. The residue is purified by chromatography (35 g silica gel, eluent: dichloromethane:methanol = 99:1).

Yield: 89.5 %, brown oil

- 313 -

b) synthesis of the final products



0.3 mmol aniline are dissolved in dichloromethane, consecutively treated with 0.33 mmol pyridine and 0.3 mmol of the p-Nitrophenylester of chloroformic acid and stirred at room temperature. After the reaction is finished, 0.33 mmol 13, suspended in dichloromethane, and 0.3 mmol DIPEA are added und and the resulting solution stirred at room temperature. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

Variant A: The reaction mixture is diluted with dichloromethane and washed three times with water. The organic phase is dried over Na₂SO₄, filtered and evaporated. The residue is recrystallized from ethylacetate.


Variant B: The reaction mixture is diluted with dichloromethane and washed three times with water. The organic phase is dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (10 g silica gel, eluent:petrol ether:ethyl acetate = 100:0 to 50:50).

Variant C: The reaction mixture is diluted with dichloromethane and washed three times with water. The organic phase is dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (10 g silica gel, eluent: dichloromethane:methanol = 100:0 to 95:5).

Variant D: The reaction mixture is treated with water, stirred and then the water phase discarded. The organic phase is washed once with water, once with brine, dried over Na₂SO₄, filtered and evaporated. The residue is taken up in hot acetonitrile. On cooling, a white precipitate is obtained, which is filtered by suction and washed with diethylether. After some time, and additional portion of product precipitated from the mother liquor which

- 314 -

is washed with diethylether/acetone (9:1). The combined products are dried in vacuo at 40 °C.





A solution of **16** in acetic acid is treated with 3 eq. sodium perborate trihydrate and heated to 55 °C. After stirring for 2 hrs at 55 °C the reaction mixture is allowed to stand overnight. Then, additional 1.5 eq. Sodium perborate trihydrate is added and the reaction mixture is stirred for another 4 hrs at 55 °C. Then the reaction mixture is cooled and poured onto ice. After multiple extraction of the reaction mixture with ethyl acetate, the combined organic phases are washed twice with 2 N sodium hydroxide solution, once with water, dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (4 g silica gel, eluent: ethyl acetate:n-heptane = 4:6 to 8:2).

Yield: 58 %, colourless crystals

Synthesis of methylene urea derivatives substituted on the methylene moiety

- 315 -



60 ml thionyl chloride are heated to 45 °C under a nitrogen atmosphere and treated slowly with 1.83 ml dimethylformamide. To this solution, 20 g pyridine-to-carbonyl acid are added in portions and the reaction mixture stirred another 45 minutes at the same temperature and then heated to 80 °C for 24 hours. The reaction mixture is evaporated and the resulting residue treated with dry toluene as a carrier and then evaporated. This procedure is repeated several times. The resulting oil is dissolved in toluene, cooled to 0 °C, slowly treated with methanol and stirred for one hour. The resulting precipitate is filtered by suction, washed with toluene and recrystallised from acetone.

Yield: 15 g (45%) of compound **17**, colourless crystals

13 g (62.5 mmol) of compound **17** are dissolved in THF together with 2.98 g (31.24 mmol) dry MgCl₂. After five minutes, 110 ml methyl amine solution (2M in THF) are added within ten minutes and the resulting suspension stirred for two hours at room temperature. The reaction mixture is treated with 120 ml water and 63 ml 1N hydrochloric acid and

- 316 -

extracted three times with ethyl acetate. The combined organic phases are washed with brine, dried over Na_2SO_4 , filtered and evaporated.

Yield: 10.5 g (98.5 %) of compound 18, colourless oil.

5 440 mg (2.58 mmol) of compound 18 are heated up together with 1.05 g (7.74 mmol) 4-Hydroxy-acetophenon in an Argon atmosphere for 18 hours. The reaction mixture is cooled, diluted with ethyl acetate, washed three times with 2N sodium hydroxide solution and twice with water, dried over Na_2SO_4 , filtered and evaporated. The residue is purified by chromatography (10 g silica gel, eluent: ethyl acetate/n-heptane = 4:6 to 7:3). Further product is isolated by crystallising the mixed fractions with petrol ether/diethyl ether.

Yield: 344 mg (49%) of compound 19, colourless solid

15 1 mg (0.37 mmol) of compound 19 is suspended in 2 ml ethanol together with 59 mg (0.37 mmol) O-Benzyl-hydroxylamine and treated with one drop concentrated sulphuric acid. The resulting clear solution is heated under reflux for 16 hours and then evaporated to dryness. The yellowish residue is treated with water, made alkaline with 1N sodium hydroxide solution (pH = 8-9) and extracted twice with ethyl acetate. The combined organic phases are dried over Na_2SO_4 , filtered and evaporated.

Yield: 151 mg (99%) of compound 20, slightly yellow oil


25 3.46 g (9.22 mmol) of compound 20 are hydrogenated at room temperature and a pressure of 5 bar in methanolic ammonium solution in the presence of Raney nickel. The reaction mixture is filtered over kieselguhr, washed with methanol and the filtrate is evaporated. The residue is purified by chromatography (120 g silica gel, eluent: $\text{CH}_2\text{Cl}_2/\text{methanol}/\text{NH}_3 = 94:6:0.1$).

30 Yield: 2.29g (91%) of compound 21, slightly yellow solid

- 317 -

Synthesis of the methylene urea derivatives substituted on the methylene moiety:

Method A




15 Variant A: The resulting precipitate is filtered by suction, washed with little dichloromethane and dried in vacuo at 40 °C.

Variant B: The reaction mixture is evaporated and the oily residue taken up in acetonitrile:water = 2:1, frozen and freeze dried overnight.

20 Variant C: The reaction mixture is evaporated to dryness and the residue dried in vacuo at 40 °C.

Method B

a) Synthesis of the trichloro aceto anilides



30 1 g substituted aniline and 1.3 eq. pyridine are dissolved in dichloromethane, cooled to 0 °C and 1.1 eq. trichloro acetic acid chloride is


slowly added. After the addition is completed, the reaction mixture is allowed warm up to room temperature and stirring is continued for 1 h. Then the reaction mixture is extracted 1M hydrochloric acid and water consecutively and the organic phase is dried over Na₂SO₄, filtered and evaporated.

R₁ = 2-OMe, 5-CF₃; colourless solid, yield: 93 %

R₁ = 3-CF₃, 4-Br; colourless solid, yield: 100 %

R₁ = 3-OCF₃; yellow solid, yield: 82 %

b) synthesis of the final products



0.15 mmol of the substituted trichloro aceto anilide and 0.15 mmol benzyl amine 5 are dissolved in DMSO, 0.15 mmol DBU are added and the mixture heated to 80 °C for 2.5-34 hrs. Depending to the respective reaction course, the working up of the reaction mixtures is done according to one of the variations given below:

Variante A: The reaction mixture is cooled, diluted with dichloromethane and washed consecutively twice with 1N hydrochloric acid and with water, dried over Na₂SO₄, filtered and evaporated. The residue is digested with a small amount of water, the resulting precipitate filtered by suction, washed with diethyl ether and dried in vacuo at 40 °C.

Variante B: The reaction mixture is cooled, diluted with dichloromethane, extracted consecutively twice with 1N hydrochloric acid and with water, dried over Na₂SO₄, filtered and evaporated. The residue is purified by chromatography (5 g silica gel, eluent: ethyl acetate/n-heptane = 4:6 to 6:4, taken up in acetonitrile: water = 2:1, frozen and freeze dried overnight.

HPLC method:

Gradient: 5.5 min; flow rate: 2.75 ml/min from 90:10 to 0:100 H₂O/ACN
Water + TFA (0.01% by vol.); acetonitrile + TFA (0.01% by vol.)

Column: Chromolith SpeedROD RP 18e 50-4.6

Wavelength: 220 nm, Rt=Retention time.

^aHPLC method:

Gradient: 9 min; flow rate: 1.5 ml/min from 80:20 to 0:100 H₂O/ACN

Water + TFA (0.01% by vol.); acetonitrile + TFA (0.01% by vol.)

Column: Lichrospher RP-select-B (5 µm/125 mm)

Wavelength: 220 nm, Rt=Retention time.

The compounds (1) to (224) as described above can preferably be produced according to the procedures described herein or in an analogous manner thereof.

The compounds (225) to (449) as described above can preferably be produced according to the procedures described herein, especially according to the ones for producing methylene urea derivatives being substituted on the methylene moiety, or in an analogous manner thereof.

The compounds (450) to (672) as described above can preferably be produced according to the procedures described herein, especially according to the ones for producing methylene urea derivatives being substituted on the methylene moiety, or in an analogous manner thereof.

Example A: Injection vials

A solution of 100 g of an active compound of the formula I and 5 g of disodium hydrogenphosphate is adjusted to pH 6.5 in 3 l of double-distilled water using 2N hydrochloric acid, sterile-filtered, dispensed into injection

- 320 -

vials, lyophilized under sterile conditions and aseptically sealed. Each injection vial contains 5 mg of active compound.

Example B: Suppositories

- 5 A mixture of 20 g of an active compound of the formula I is fused with 100 g of soya lecithin and 1400 g of cocoa butter, poured into moulds and allowed to cool. Each suppository contains 20 mg of active compound.

Example C: Solution

- 10 A solution of 1 g of an active compound of the formula I, 9.38 g of $\text{NaH}_2\text{PO}_4 \cdot 2 \text{H}_2\text{O}$, 28.48 g of $\text{Na}_2\text{HPO}_4 \cdot 12 \text{H}_2\text{O}$ and 0.1 g of benzalkonium chloride in 940 ml of double-distilled water is prepared. It is adjusted to pH 6.8, made up to 1 l and sterilized by irradiation. This solution can be used in the form of eye drops.

Example D: Ointment

- 500 mg of an active compound of the formula I is mixed with 99.5 g of petroleum jelly under aseptic conditions.

Example E: Tablets

- 25 A mixture of 1 kg of active compound of the formula I, 4 kg of lactose, 1.2 kg of potato starch, 0.2 kg of talc and 0.1 kg of magnesium stearate is compressed to give tablets in a customary manner such that each tablet contains 10 mg of active compound.

Example F: Coated tablets

30

- 321 -

Analogously to Example E, tablets are pressed and are then coated in a customary manner using a coating of sucrose, potato starch, talc, tragacanth and colourant.

Example G: Capsules

- 5 2 kg of active compound of the formula I are dispensed into hard gelatin capsules in a customary manner such that each capsule contains 20 mg of the active compound.

Example H: Ampoules

- 10 A solution of 1 kg of active compound of the formula I in 60 l of double-distilled water is sterile-filtered, dispensed into ampoules, lyophilized under sterile conditions and aseptically sealed. Each ampoule contains 10 mg of active compound.

20

25

30

- 322 -

Claims

1. Methylene urea derivatives of formula I

5

A-D-B

(I)

wherein

D is a bivalent methylene urea moiety, or a derivative thereof,

10

A is a unsubstituted or substituted moiety of up to 40 carbon atoms of the formula: $-L-(M-L')_n$, where L is a 5, 6 or 7 membered cyclic structure, preferably selected from the group consisting of aryl, heteroaryl, arylene and heteroarylene, bound directly to D, L' comprises an optionally substituted cyclic moiety having at least 5 members, preferably selected from the group consisting of aryl, heteroaryl, alkyl, cycloalkyl and heterocyclyl, M is a bond or a bridging group having at least to one atom, α is an integer of from 1-4; and each cyclic structure of L and L' contains 0-4 members of the group consisting of nitrogen, oxygen and sulfur, wherein L' is preferably substituted by at least one substituent selected from the group consisting of -

20

 SO_2R_x , $-C(O)R_x$ and $-C(NR_y)R_z$

25

B is a substituted or unsubstituted, up to tricyclic aryl or heteroaryl moiety of up to 30 carbon atoms, preferably of up to 20 carbon atoms, comprising at least one 5-, 6-, or 7-membered cyclic structure, preferably a 5- or 6-membered cyclic structure, bound directly to D containing 0-4 members of the group consisting of nitrogen, oxygen and sulfur, wherein said cyclic structure directly bound to D is preferably selected from the group

30

- 323 -

consisting of aryl, heteroaryl and heterocyclyl, R_y is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally halosubstituted, up to per halo,

5

R_z is hydrogen or a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen;

10

R_x is R_z or NR_aR_b , where R_a and R_b are

15

a) independently hydrogen, a carbon based moiety of up to 30 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen, or

20

$-OSi(R_i)_3$ where R_i is hydrogen or a carbon based moiety of up to 24 carbon atoms optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, hydroxy and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen;

or

30

- 324 -

b) R_a and R_b together form a 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O, or a substituted 5-7 member heterocyclic structure of 1-3 heteroatoms selected from N, S and O substituted by halogen, hydroxy or carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen; or

c) one of R_a or R_b is $-C(O)-$, a C_1-C_3 divalent alkylene group or a substituted C_1-C_3 divalent alkylene group bound to the moiety L to form a cyclic structure with at least 5 members, wherein the substituents of the substituted C_1-C_3 divalent alkylene group are selected from the group consisting of halogen, hydroxy, and carbon based substituents of up to 24 carbon atoms, which optionally contain heteroatoms selected from N, S and O and are optionally substituted by halogen;

where B is substituted, L is substituted or L' is additionally substituted, the substituents are selected from the group consisting of halogen, up to per-halo, and W_x , where x is 0-3;

wherein each W is independently selected from the group consisting of $-CN$, $-CO_2R$, $-C(O)NR^5R^5$, $-C(O)R^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-NR^5R^5$, $-NR^5C(O)OR^5$, $-NR^5C(O)R^5$, $-Q-A$, and carbon based moieties of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O and optionally substituted by one or more substituents independently selected from the groups consisting of $-CN$, $-CO_2R$, $-C(O)NR^5R^5$, $-C(O)R^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-NR^5R^5$, $-NR^5C(O)OR^5$, $-NR^5C(O)R^5$ and halogen up to per-halo;

- 325 -

with each R^5 independently selected from H or a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O and optionally substituted by halogen, wherein Q is $-O-$, $-S-$, $-N(R^5)-$, $-(CH_2)_\beta-$, $-C(O)-$, $-CH(OH)-$, $-(CH_2)_\beta-S-$, $-(CH_2)_\beta N(R^5)-$, $-O(CH_2)_\beta-CHHal-$, $-CHal-$, $-S-(CH_2)_\beta-$ and $-N(R^5)(CH_2)_\beta-$ where $\beta = 1-3$, and Hal is halogen; and Ar is 5- or 6-member aromatic structure containing 0-2 members selected from the group consisting of nitrogen, oxygen and sulfur, which is optionally substituted by halogen, up to per-halo, and optionally substituted by Z $\delta 1$

wherein $\delta 1$ is 0 to 3 and each Z is independently selected from the group consisting of $-CN$, $-CO_2R^5$, $-C(O)NR^5R^5$, $-C(O)R^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-SO_2R^5$, $-SO_3H$, $-NR^5R^5$, $-NR^5C(O)OR^5$, $-NR^5C(O)R^5$, and a carbon based moiety of up to 24 carbon atoms, optionally containing heteroatoms selected from N, S and O and optionally substituted by one or more substituents selected from the group consisting of $-CN$, $-CO_2R^5$, $-C(O)NR^5R^5$, $-C(O)R^5$, $-NO_2$, $-OR^5$, $-SR^5$, $-SO_2R^5$, $-SO_3H$, $-NR^5R^5$, $-NR^5C(O)OR^5$, $-NR^5C(O)R^5$, and the pharmaceutically acceptable derivatives, salts and solvates thereof.

2. Methylene urea derivative according to claim 1, characterised in that each M independently from one another represents a bond OR is a bridging group, selected from the group consisting of $(CR^5R^5)_n$, or $(CHR^5)_n-Q-(CHR^5)_n$, wherein

Q is selected from a group consisting of O, S, $N-R^5$, $(CHal)_n$, $(O-CHR^5)_n$, $(CHR^5-O)_n$, $CR^5=CR^5$, $(O-CHR^5CHR^5)_n$, $(CHR^5CHR^5-O)_n$, $C=O$, $C=S$, $C=NR^5$, $CH(OR^5)$, $C(OR^5)(OR^5)$,

- 326 -


$C(=O)O$, $OC(=O)$, $OC(=O)O$, $C(=O)N(R^5)C(=O)$, $OC(=O)N(R^5)$, $N(R^5)C(=O)O$, $CH=N-NR^5$, $OC(O)NR^5$, $NR^5C(O)O$, $S=O$, SO_2 , SO_2NR^5 und NR^5SO_2 , wherein

5 R^5 is in each case independently selected from the meanings given above, preferably hydrogen, halogen, alkyl, aryl, aralkyl,

h, i are independently from each other 0, 1, 2, 3, 4, 5, or 6, preferably 0, 1, 2 or 3, and

10 j is 1, 2, 3, 4, 5 or 6, preferably 0, 1, 2 or 3.

3. Methylene urea derivative according to claim 1 or 2, selected from the compounds of formula II,



20 wherein

Ar^1, Ar^2 are selected independently from one another from aromatic hydrocarbons containing 6 to 14 carbon atoms and ethylenical unsaturated or aromatic heterocyclic residues containing 3 to 10 carbon atoms and one or two hetero atoms, independently selected from N, O und S,

30 R^6, R^7 are independently selected from a the meanings given for R^6, R^9 and R^{10} , or

- 327 -

R^6 and R^7 together form a carbocyclic residue comprising 3 to 7 carbon atoms or a heterocyclic residue comprising 1, 2 or 3 hetero atoms, selected from the group consisting of O, N and S, and 2 to 6 carbon atoms, said carbocyclic or heterocyclic residue being unsubstituted or comprising 1, 2 or 3 substituents, selected from the meanings given for R^6, R^9 and R^{10} ,

E, G, M, Q and U are selected, independently from one another, from carbon atoms and nitrogen atoms, with the proviso that one or more of E, G, M, Q and U are carbon atoms and that X is bonded to a carbon atom,

R^8, R^9 and R^{10} are independently selected from a group consisting of

15 H, A, cycloalkyl comprising 3 to 7 carbon atoms, Hal, CH_2Hal , $CH(Hal)_2$, $C(Hal)_3$, NO_2 , $(CH_2)_hCN$, $(CH_2)_hNR^{11}R^{12}$, $(CH_2)_hO(CH_2)_kNR^{11}R^{12}$, $(CH_2)_hNR^{11}(CH_2)_kNR^{11}R^{12}$, $(CH_2)_hO(CH_2)_kOR^{11}$, $(CH_2)_hNR^{11}(CH_2)_kOR^{12}$, $(CH_2)_hCOOR^{13}$, $(CH_2)_hCOR^{13}$, $(CH_2)_hCONR^{11}R^{12}$, $(CH_2)_hNR^{11}COR^{13}$, $(CH_2)_hNR^9CONR^{11}R^{12}$, $(CH_2)_hNR^{11}SO_2A$, $(CH_2)_hSO_2NR^{11}R^{12}$, $(CH_2)_hS(O)_hR^{13}$, $(CH_2)_hOC(O)R^{13}$, $(CH_2)_hCOR^{13}$, $(CH_2)_hSR^{11}$, $CH=N-OA$, $CH_2CH=N-OA$, $(CH_2)_hNHOA$, $(CH_2)_hCH=N-N^{11}$, $(CH_2)_hOC(O)NR^{11}R^{12}$, $(CH_2)_hNR^{11}COOR^{13}$, $(CH_2)_hN(R^{11})CH_2CH_2OR^{13}$, $(CH_2)_hN(R^{11})CH_2CH_2OCF_3$, $(CH_2)_hN(R^{11})C(R^{13})HCOOR^8$, $(CH_2)_hN(R^{11})$, $C(R^{13})HCOOR^8$, $(CH_2)_hN(R^{11})CH_2CH_2N(R^{12})CH_2COOR^8$, $(CH_2)_hN(R^9)CH_2CH_2NR^{12}R^8$, $CH=CHCOOR^{13}$, $CH=CHCH_2NR^{11}R^{12}$, $CH=CHCH_2NR^{11}R^{12}$, $CH=CHCH_2OR^{13}$, $(CH_2)_hN(COOR^{13})COOR^{14}$,

20

25

30

- 328 -

5 $(CH_2)_mN(CONH_2)COOR^{13}$, $(CH_2)_mN(CONH_2)CONH_2$,
 $(CH_2)_mN(CH_2COOR^{13})COOR^{14}$,
 $(CH_2)_mN(CH_2CONH_2)COOR^{13}$,
 $(CH_2)_mN(CH_2CONH_2)CONH_2$, $(CH_2)_mCHR^{13}COR^{14}$,
 $(CH_2)_mCHR^{13}COOR^{14}$, $(CH_2)_mCHR^{13}CH_2OR^{14}$,
 $(CH_2)_mOCN$ and $(CH_2)_mNCO$, wherein

10 R^{11} , R^{12}
are independently selected from a group consisting of
H, A, $(CH_2)_mAr^3$ and $(CH_2)_mHet$, or in $NR^{11}R^{12}$,

form, together with the N-atom they are bound to, a 5-,
6- or 7- membered heterocyclyus which optionally
contains 1 or 2 additional hetero atoms, selected from
N, O and S,

15 R^{13} , R^{14}
are independently selected from a group consisting of
H, Hal, A, $(CH_2)_mAr^4$ and $(CH_2)_mHet$,

20 A
is selected from the group consisting of alkyl, alkenyl,
cycloalkyl, alkylenecycloalkyl, alkoxy, alkoxyalkyl and
saturated heterocyclyl,

25 Ar^3 , Ar^4
are independently from one another aromatic hydrocarbon
residues comprising 5 to 12 and preferably 5 to 10 carbon
atoms which are optionally substituted by one or more
substituents, selected from a group consisting of A, Hal,
 NO_2 , CN, OR¹⁵, $NR^{15}R^{16}$, COOR¹⁵, CONR¹⁵R¹⁶,
 $NR^{15}COR^{16}$, $NR^{15}CONR^{15}R^{16}$, $NR^{16}SO_2A$, COR¹⁵,
 $SO_2R^{15}R^{16}$, $S(O)_2A$ and OOCR¹⁵,

30 Het
is a saturated, unsaturated or aromatic heterocyclic
residue which is optionally substituted by one or more

- 329 -

substituents, selected from a group consisting of A, Hal,
 NO_2 , CN, OR¹⁵, $NR^{15}R^{16}$, COOR¹⁵, CONR¹⁵R¹⁶,
 $NR^{15}COR^{16}$, $NR^{15}CONR^{15}R^{16}$, $NR^{16}SO_2A$, COR¹⁵,
 $SO_2R^{15}R^{16}$, $S(O)_2A$ and OOCR¹⁵,

5 R^{15} , R^{16}
are independently selected from a group consisting of H,
A, and $(CH_2)_mAr^6$, wherein

10 Ar^6
is a 5- or 6-membered aromatic hydrocarbon which is
optionally substituted by one or more substituents selected
from a group consisting of methyl, ethyl, propyl, 2-propyl,
tert-butyl, Hal, CN, OH, NH₂ and CF₃.

k , n and m are independently of one another 0, 1, 2, 3, 4, or 5;

15 X
represents a bond or is $(CR^{11}R^{12})_m$, or
 $(CHR^{11})_m-Q-(CHR^{12})_n$, wherein

20 Q
is selected from a group consisting of O, S, N-R¹⁵, $(CH_2)_2$,
 $(O-CHR^{18})_n$, $(CHR^{18}-O)_n$, $CR^{18}=CR^{19}$, $(O-CHR^{18}CHR^{19})_n$,
 $CHR^{18}CHR^{19}-O)_n$, C=O, C=S, C=NR¹⁵, $CH(OR^{15})$,
 $C(OR^{15})(OR^{20})$, C(=O)O, OC(=O), OC(=O)O, C(=N)(R¹⁵),
 $N(R^{15})C(=O)$, OC(=O)N(R¹⁵), $N(R^{15})C(=O)O$, CH=N-O,
CH=N-NR¹⁵, OC(O)NR¹⁵, $NR^{15}C(O)O$, S=O, SO₂, SO₂NR¹⁵
and NR¹⁵SO₂, wherein

25 h , l
are independently from each other 0, 1, 2, 3, 4, 5 or 6, and

J
is 1, 2, 3, 4, 5 or 6,

30 Y
is selected from O, S, NR²¹, $C(R^{22})_2NO_2$, $C(R^{22})_2CN$ and
 $C(CN)_2$, wherein

-330-

R^{21} is independently selected from the meanings given for R^{13} , R^{14} , and

R^{22} is independently selected from the meanings given for R^{11} , R^{12} , R^{13} , R^{14} , R^{15} , R^{16} , R^{17} , R^{18} , R^{19} , R^{20} , R^{21} , R^{23} , R^{24} , R^{25} , R^{26} , R^{27} , R^{28} , R^{29} , R^{30} , R^{31} , R^{32} , R^{33} , R^{34} , R^{35} , R^{36} , R^{37} , R^{38} , R^{39} , R^{40} , R^{41} , R^{42} , R^{43} , R^{44} , R^{45} , R^{46} , R^{47} , R^{48} , R^{49} , R^{50} , R^{51} , R^{52} , R^{53} , R^{54} , R^{55} , R^{56} , R^{57} , R^{58} , R^{59} , R^{60} , R^{61} , R^{62} , R^{63} , R^{64} , R^{65} , R^{66} , R^{67} , R^{68} , R^{69} , R^{70} , R^{71} , R^{72} , R^{73} , R^{74} , R^{75} , R^{76} , R^{77} , R^{78} , R^{79} , R^{80} , R^{81} , R^{82} , R^{83} , R^{84} , R^{85} , R^{86} , R^{87} , R^{88} , R^{89} , R^{90} , R^{91} , R^{92} , R^{93} , R^{94} , R^{95} , R^{96} , R^{97} , R^{98} , R^{99} , R^{100} , R^{101} , R^{102} , R^{103} , R^{104} , R^{105} , R^{106} , R^{107} , R^{108} , R^{109} , R^{110} , R^{111} , R^{112} , R^{113} , R^{114} , R^{115} , R^{116} , R^{117} , R^{118} , R^{119} , R^{120} , R^{121} , R^{122} , R^{123} , R^{124} , R^{125} , R^{126} , R^{127} , R^{128} , R^{129} , R^{130} , R^{131} , R^{132} , R^{133} , R^{134} , R^{135} , R^{136} , R^{137} , R^{138} , R^{139} , R^{140} , R^{141} , R^{142} , R^{143} , R^{144} , R^{145} , R^{146} , R^{147} , R^{148} , R^{149} , R^{150} , R^{151} , R^{152} , R^{153} , R^{154} , R^{155} , R^{156} , R^{157} , R^{158} , R^{159} , R^{160} , R^{161} , R^{162} , R^{163} , R^{164} , R^{165} , R^{166} , R^{167} , R^{168} , R^{169} , R^{170} , R^{171} , R^{172} , R^{173} , R^{174} , R^{175} , R^{176} , R^{177} , R^{178} , R^{179} , R^{180} , R^{181} , R^{182} , R^{183} , R^{184} , R^{185} , R^{186} , R^{187} , R^{188} , R^{189} , R^{190} , R^{191} , R^{192} , R^{193} , R^{194} , R^{195} , R^{196} , R^{197} , R^{198} , R^{199} , R^{200} , R^{201} , R^{202} , R^{203} , R^{204} , R^{205} , R^{206} , R^{207} , R^{208} , R^{209} , R^{210} , R^{211} , R^{212} , R^{213} , R^{214} , R^{215} , R^{216} , R^{217} , R^{218} , R^{219} , R^{220} , R^{221} , R^{222} , R^{223} , R^{224} , R^{225} , R^{226} , R^{227} , R^{228} , R^{229} , R^{230} , R^{231} , R^{232} , R^{233} , R^{234} , R^{235} , R^{236} , R^{237} , R^{238} , R^{239} , R^{240} , R^{241} , R^{242} , R^{243} , R^{244} , R^{245} , R^{246} , R^{247} , R^{248} , R^{249} , R^{250} , R^{251} , R^{252} , R^{253} , R^{254} , R^{255} , R^{256} , R^{257} , R^{258} , R^{259} , R^{260} , R^{261} , R^{262} , R^{263} , R^{264} , R^{265} , R^{266} , R^{267} , R^{268} , R^{269} , R^{270} , R^{271} , R^{272} , R^{273} , R^{274} , R^{275} , R^{276} , R^{277} , R^{278} , R^{279} , R^{280} , R^{281} , R^{282} , R^{283} , R^{284} , R^{285} , R^{286} , R^{287} , R^{288} , R^{289} , R^{290} , R^{291} , R^{292} , R^{293} , R^{294} , R^{295} , R^{296} , R^{297} , R^{298} , R^{299} , R^{300} , R^{301} , R^{302} , R^{303} , R^{304} , R^{305} , R^{306} , R^{307} , R^{308} , R^{309} , R^{310} , R^{311} , R^{312} , R^{313} , R^{314} , R^{315} , R^{316} , R^{317} , R^{318} , R^{319} , R^{320} , R^{321} , R^{322} , R^{323} , R^{324} , R^{325} , R^{326} , R^{327} , R^{328} , R^{329} , R^{330} , R^{331} , R^{332} , R^{333} , R^{334} , R^{335} , R^{336} , R^{337} , R^{338} , R^{339} , R^{340} , R^{341} , R^{342} , R^{343} , R^{344} , R^{345} , R^{346} , R^{347} , R^{348} , R^{349} , R^{350} , R^{351} , R^{352} , R^{353} , R^{354} , R^{355} , R^{356} , R^{357} , R^{358} , R^{359} , R^{360} , R^{361} , R^{362} , R^{363} , R^{364} , R^{365} , R^{366} , R^{367} , R^{368} , R^{369} , R^{370} , R^{371} , R^{372} , R^{373} , R^{374} , R^{375} , R^{376} , R^{377} , R^{378} , R^{379} , R^{380} , R^{381} , R^{382} , R^{383} , R^{384} , R^{385} , R^{386} , R^{387} , R^{388} , R^{389} , R^{390} , R^{391} , R^{392} , R^{393} , R^{394} , R^{395} , R^{396} , R^{397} , R^{398} , R^{399} , R^{400} , R^{401} , R^{402} , R^{403} , R^{404} , R^{405} , R^{406} , R^{407} , R^{408} , R^{409} , R^{410} , R^{411} , R^{412} , R^{413} , R^{414} , R^{415} , R^{416} , R^{417} , R^{418} , R^{419} , R^{420} , R^{421} , R^{422} , R^{423} , R^{424} , R^{425} , R^{426} , R^{427} , R^{42

p, r are independently from one another 0, 1, 2, 3, 4 or 5,

q is 0, 1, 2, 3 or 4, preferably 0, 1 or 2,


u is 0, 1, 2 or 3, preferably 0, 1 or 2,

and

Hal is independently selected from a group consisting of F, Cl, Br and I;


and the pharmaceutically acceptable derivatives, salts and solvates thereof.

4. Methylene urea derivative according to one of the claims 1 to 3, selected from the compounds of formula Iic, IId, IIf, IIg, IIh, IIi, IIj, IIk, IIl, IIm, IIo, IIp, IIq, IIr, IIs, IIt, IIu, IIv, IIw and IIx,




30


-331-




5




10




15



20








25









30


- 332 -




30

- 333 -







5



10



15



20

wherein R^6 , R^7 , R^8 , p , Ar^1 , Y , X , R^9 and q are as defined in claim 3, R^{10} is H or as defined in claim 3; and the pharmaceutically acceptable derivatives, salts and solvates thereof.

25

5. Methylene urea derivative according to claim one of the claims 1, 2 or 3, selected from the compounds (1) to (224) of table 1, the

30

compounds (225) to (449) of table 2 and/or the compounds (450) to (672) of table 3, and the pharmaceutically acceptable derivatives, salts and solvates thereof.

6. Methylene urea derivative according to claim one of the claims 1, 2 or 3, selected from the compounds (673) to (758), the compounds (759) to (825) and/or the compounds (826) to (874); and the pharmaceutically acceptable derivatives, salts and solvates thereof.

5

7. Methylene urea derivative according to one of the claims 1 to 6 as a medicament.

10

8. Methylene urea derivative according to one of the claims 1 to 6 as a kinase inhibitor.

15

9. Methylene urea derivative according to claim 8, characterized in that the kinases are selected from raf-kinases.

15

10. Pharmaceutical composition, characterised in that it contains one or more compounds according to one of the claims 1 to 6.

20

11. Pharmaceutical composition according to claim 10, characterised in that it contains one or more additional compounds, selected from the group consisting of physiologically acceptable excipients, auxiliaries, adjuvants, carriers and pharmaceutical active ingredients other than the compounds according to one of the claims 1 to 6.

25

12. Process for the manufacture of a pharmaceutical composition, characterised in that one or more compounds according to one of the claims 1 to 6 and one or more compounds, selected from the group consisting of carriers, excipients, auxiliaries and pharmaceutical

30

- 336 -

active ingredients other than the compounds according to one of the claims 1 to 6, is processed by mechanical means into a pharmaceutical composition that is suitable as dosageform for application and/or administration to a patient.

- 5 13. Use of a compound according to one of the claims 1 to 6 as a pharmaceutical.

- 10 14. Use of a compound according to one of the claims 1 to 6 in the treatment and/or prophylaxis of disorders.

- 15 15. Use of a compound according to one of the claims 1 to 6 for producing a pharmaceutical composition for the treatment and/or prophylaxis of disorders.

16. Use according to claim 14 or 15, characterised in that the disorders are caused, mediated and/or propagated by raf-kinases.

- 20 17. Use according to claim 14, 15 or 16, characterised in that the disorders are selected from the group consisting of hyperproliferative and nonhyperproliferative disorders.

- 25 18. Use according to claim 14, 15, 16 or 17, characterised in that the disorder is cancer.

19. Use according to claim 14, 15, 16 or 17, characterised in that the disorder is noncancerous.

- 30 20. Use according to claim 14, 15, 16, 17 or 19, characterised in that the disorders are selected from the group consisting of psoriasis, arthritis, inflammation, endometriosis, scarring, Helicobacter pylori infection,

- 337 -

Influenza A, benign prostatic hyperplasia, immunological diseases, autoimmune diseases and immunodeficiency diseases.

- 5 21. Use according to one of the claims 14 to 18, characterised in that the disorders are selected from the group consisting of melanoma, brain cancer, lung cancer, squamous cell cancer, bladder cancer, gastric cancer, pancreatic cancer, hepatic cancer, renal cancer, colorectal cancer, breast cancer, head cancer, neck cancer, oesophageal cancer, gynaecological cancer, ovarian cancer, ovary cancer, uterine cancer, prostate cancer, thyroid cancer, lymphoma, chronic leukaemia and acute leukaemia.
- 10

- 15 22. Use according to one of the claims 14 to 19, characterised in that the disorders are selected from the group consisting of arthritis, restenosis, fibrotic disorders, mesangial cell proliferative disorders, diabetic nephropathy, malignant nephrosclerosis, thrombotic microangiopathy syndromes, organ transplant rejection, glomerulopathies, metabolic disorders, inflammation, solid tumors, rheumatic arthritis, diabetic retinopathy, and neurodegenerative diseases.
- 20

23. Use according to one of the claims 14 to 17, characterised in that the disorders are selected from the group consisting of rheumatoid arthritis, inflammation, autoimmune disease, chronic obstructive pulmonary disease, asthma, inflammatory bowel disease, fibrosis, atherosclerosis, restenosis, vascular disease, cardiovascular disease, inflammation, renal disease and angiogenesis disorders.
- 25

- 30 24. Use of a compound according to one of the claims 1 to 6 as a raf-kinase inhibitor.

25. Use according to claim 24, characterised in that the raf-kinase is selected from the group consisting of A-Raf, B-Raf and c-Raf1.

26. Method for the treatment and/or prophylaxis of disorders,

characterised in that one or more compounds according to one of the claims 1 to 6 is administered to a patient in need of such a treatment.

27. Method according to claim 26, characterised in that the one or more compounds according to one of the claims claim 1 to 6 are administered as a pharmaceutical composition according to claim 10 or 11.

28. Method for the treatment and/or prophylaxis of disorders according to claim 27, characterised in that the disorders are as defined in one of the claims 16 to 23.

29. Method for the treatment according to claim 28, characterised in that the disorder is cancerous cell growth mediated by raf-kinase.

30. Method for producing compounds of formula II, characterised in that

a) a compound of formula III



wherein


FG is a functional group, selected from

$$-\text{N}=\text{C}=\text{Y} \text{ and } -\text{NH}-(\text{C}=\text{Y})-\text{LG},$$

wherein Y is as defined as in claim 3 and LG is a leaving group,

is reacted

b) with a compound of IV,



wherein

L^2, L^3 are independently from one another H or a metal ion, and $R^6, R^7, E, G, M, Q, U, R^9, q, X, Ar^2, R^{10}$ and r are as defined in claim 3.

and optionally

c) isolating and/or treating the compound of formula II obtained by said reaction with an acid, to obtain the salt thereof

31. Compound of formula III,




wherein

FG is a functional group, selected from $-\text{N}=\text{C}=\text{Y}$ and $-\text{NH}-(\text{C}=\text{Y})-\text{LG}$,

- 340 -

wherein Y is as defined as in claim 3 and LG is a leaving group.

32. Compound of formula IV,



10 wherein

15 L^2, L^3 are independently from one another H or a metal ion, and $R^6, R^7, E, G, M, Q, U, R^8, q, X, Ar^2, R^{10}$ and r are as defined in claim 3.

25

30

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization

International Bureau



(10) International Publication Number

(43) International Publication Date
6 May 2004 (06.05.2004)

PCT

WO 2004/037789 A3

(51) International Patent Classification: C07D 213/68, (74) Common Representative: MERCK PATENT GMBH; 213/75, 401/12, 413/12, 409/12, 417/12, A61K 31/44, Frankfurter Strasse 250, 64293 Darmstadt (DE); 31/4427, A61P 29/00

(21) International Application Number: PCT/EP2003/011134

(22) International Filing Date: 8 October 2003 (08.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 24 October 2002 (24.10.2002) EP 02023906.7
28 July 2003 (28.07.2003) US 60/490,285

(71) Applicant (for all designated States except US): MERCK PATENT GMBH (DE/DE); Frankfurter Strasse 250, 64293 Darmstadt (DE);

(72) Inventors, and (73) Inventors/Applicants (for US only): BUCHSTALLER, Hans-Peter (AT/DE); Heinrichstrasse 54, 64331 Weiterstadt (DE); WIESNER, Matthias (DE/DE); Beethovenring 10, 64342 Seelheim-Jugenheim (DE); SCHAUT, Oliver (DE/DE); Eschenstrasse 32, 63517 Rodebach (DE); AMENDT, Christiane (DE/DE); Burkhausstrasse 22, 64289 Darmstadt (DE); ZENKE, Frank (DE/DE); Schulzengasse 7, 64291 Darmstadt (DE); SIRENBERG, Christian (DE/DE); Teannustrasse 10, 64289 Darmstadt (DE); GRELL, Matthias (DE/DE); Lindeweg 44, 64291 Darmstadt (DE); FINSINGER, Dirk (DE/DE); Im Fiedlersee 5, 64291 Darmstadt (DE).

Published: — with international search report — before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(86) Date of publication of the International search report: 28 October 2004

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2004/037789 A3

(54) Title: METHYLENE UREA DERIVATIVES AS RAH-KINASE INHIBITORS

(57) Abstract: The present invention relates to methylene urea derivatives of formula (I), the use of the compounds of formula (I) as inhibitors of raf-kinase, the use of the compounds of formula (I) for the manufacture of a pharmaceutical composition and a method of treatment, comprising administering said pharmaceutical composition to a patient.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/11134

A. CLASSIFICATION OF SUBJECT MATTER IPC 7 C07D213/68 C07D213/75 C07D401/12 C07D413/12 C07D409/12 C07D417/12 A61K31/44 A61K31/4427 A61P29/00	
According to International Patent Classification (IPC) or to both national classification and IPC B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D	
Documentation searched other than minimum documentation to the extent that such documents are included in this field searched	
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, CHEM ABS Data, BEILSTEIN Data, WPI Data	
C. DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	Citation of document, with indication, where appropriate, of the relevant passages
X	WO 01/38324 A (LOVELL PETER JOHN ; DEAN DAVID KENNETH (GB) ; SMITHKLINE BEECHAM PLC (G) 31 May 2001 (2001-05-31) see examples 78-81 and claims 1, 11-15
Y	
X	WO 02/24679 A (SHIMADA MITSUYUKI ; BAYER AG (DE) ; ZIEGELBAUER KARL B (DE) ; KORIYAMA Y) 28 March 2002 (2002-03-28) see e.g. example 32-14 and claims 1-4, 6, 7, 9-15 ----- -/-
	Referred to claim No. 1-3, 7-32 1-32 1-3, 7, 8, 10-15, 17-23, 26-28, 30-32
X Further documents are listed in the continuation of box C. X Patent family members are listed in annex.	
* Special categories of cited documents : *X* document defining the general state of the art which is not considered to be of particular relevance *Y* earlier document but published on or after the international filing date *Z* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another claim or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	
Date of the actual compilation of the international search	
Date of mailing of the international search report	
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentkan 2 L-2280 HV Rixwijk Tel. (+31) - (0) 43 20 30 30 Fax (+31) - (0) 43 20 30 30	23 July 2004 18.08.2004 Traegler-Goe[de], M Authorized officer

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/11134

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	
X	US 4 057 636 A (PETERSEN HANS JORGENSEN) 8 November 1977 (1977-11-08) see the 5 benzylderivatives of table A and examples 37 and 57, its production and pharmacological activity	1,2,7, 10-15, 19,23, 26-28,31
X	US 5 441 984 A (HEATH, WILLIAM F., JR. ET AL) 15 August 1995 (1995-08-15) see examples 6 and 10, its production and activity	1,2,7, 10-15, 19,23, 26-28,31
X,P	HOPKINS, THUTAM P. ET AL: "Solid-Phase Synthesis of Trisubstituted Guanidines" JOURNAL OF COMBINATORIAL CHEMISTRY, 4(2), 167-174 CODEN: JCCHFF; ISSN: 1520-4766, 2002, XP002289534 see compound with RN 409080-96-2 and its production	1-3, 30-32
X	WO 00/61561 A1 (SHIONOGI BIORESEARCH CORP., USA) 19 October 2000 (2000-10-19) see inter alia many especially example 26, its production and activity	1,2,7, 10-15, 17,18, 21,22, 26-28, 30-32
X	WO 00/61559 A1 (SHIONOGI RESEARCH CORP., USA) 19 October 2000 (2000-10-19) see inter alia many especially example 26, its production and activity	1-3,7, 10-15, 17,18, 21,22, 26-28
X	see e.g. examples 14-16 and scheme on p. 16 WO 95/18126 A1 (FUJISAWA PHARMACEUTICAL CO., LTD., JAPAN) 6 July 1995 (1995-07-06) see compound with CAS RN 168971-19-5	1,2,7, 10-15, 19,20, 22,23, 26-28
X	WO 02/02534 A1 (ASTRAZENECA AB, SWED.; ASTRAZENECA UK LTD.) 10 January 2002 (2002-01-10) see examples 6, 16 and 31 -/-	1,2,7,8, 10-15, 17,18, 21,22, 26-28

Form PCT/ISA/WFO (continuation of second sheet) (January 2004)

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP 03/11134

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the relevant passages	
X	WO 01/57008 A1 (BASF AKTIENGESELLSCHAFT, GERMANY) 9 August 2001 (2001-08-09) see e.g. inter alia examples 24 and 26-31 and in combination with scheme II	1,2,7,8, 10-15, 17,18, 20-22, 26-28,31

Form PCT/ISA/WFO (continuation of second sheet) (January 2004)

INTERNATIONAL SEARCH REPORT

International application No.
PCT/EP 03/11134

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☒ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
1-3, 7-32(part)
see FURTHER INFORMATION sheet PCT/ISA/210
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see additional sheet

1. ☒ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in this claim; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☒ No protest accompanied the payment of additional search fees.

International Application No. PCT/EP 03/11134

FURTHER INFORMATION CONTINUED FROM PCT/ISA/210

Continuation of Box I.2

Claims Nos.: 1-3, 7-32(part)

Present claims 1 to 3, 7 to 32 relate to an extremely large number of possible compounds, processes for their production, compositions and methods. Support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT is to be found, however, for only a very small proportion of the compounds claimed. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible.

Furthermore, the initial phase of the search revealed a very large number of documents relevant to the issue of novelty. So many documents were retrieved that it is impossible to determine which parts of the claim(s) may be said to define subject-matter for which protection might legitimately be sought (Article 6 PCT).

For these two reasons, a meaningful search over the whole breadth of the claims 1 to 3 and 7 to 32 is impossible. Consequently, the search has been carried out for those parts of the claims which appear to be supported and disclosed, namely the search has been restricted to the compounds of claim 4 to 6 and the claims depending thereon, but only as far as the specified compounds of claims 4, 5 and 6 are concerned.

The applicant's attention is drawn to the fact that claims relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure. If the application proceeds into the regional phase before the EPO, the applicant is reminded that a search may be carried out during examination before the EPO (see EPO guideline C-VI, 8.5), should the problems which led to the Article 17(2) declaration be overcome.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 03/11134

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0138324	A	31-05-2001	
AU	1529901 A	04-06-2001	
AU	1623601 A	04-06-2001	
BR	0015532 A	25-06-2002	
CA	2395564 A1	31-05-2001	
CZ	20021746 A3	16-10-2002	
EP	1232153 A2	21-08-2002	
EP	1233769 A1	28-08-2002	
EP	0138324 A2	31-05-2001	
HU	0203403 A2	28-02-2003	
JP	2003514860 T	22-04-2003	
JP	2003514906 T	22-04-2003	
NO	20022318 A	15-05-2002	
NZ	518032 A	27-02-2004	
PL	355912 A1	31-05-2004	
TR	200201364 T2	21-10-2002	
WO	0137835 A1	31-05-2001	
ZA	200203987 A	25-11-2003	
WO 0224679	A	28-03-2002	
JP	2002114777 A	16-04-2002	
AU	8987301 A	02-04-2002	
AU	8987501 A	02-04-2002	
AU	9185201 A	02-04-2002	
B6	107632 A	28-11-2003	
BR	0114073 A	17-06-2003	
CA	2422923 A1	28-03-2002	
CN	1476442 T	18-02-2004	
WO	0224679 A1	28-03-2002	
WO	0224651 A1	28-03-2002	
EP	0224693 A1	16-07-2003	
EP	1326856 A1	29-12-2003	
HU	0303124 A2	01-07-2003	
MA	25836 A1	21-03-2003	
NO	20031307 A	08-01-2004	
SK	3452003 A3	13-05-2003	
US	6562811 B1		
WO 02062750	A	15-08-2002	
BR	0206955 A	09-03-2004	
CA	2436659 A1	15-08-2002	
CZ	20032122 A3	15-10-2003	
EP	1368308 A1	10-12-2003	
HU	0304060 A2	28-04-2004	
NO	20033505 A	07-10-2003	
SK	9952003 A3	08-01-2004	
WO	02062750 A1	15-08-2002	
US	2003096844 A1	22-05-2003	
US	2003232859 A1	18-12-2003	
EP 0839803	A	06-05-1998	
CA	2218360 A1	30-04-1998	
CN	1181377 A, B	13-05-1998	
EP	0839803 A1	06-05-1998	
JP	10182588 A	07-07-1998	
KR	272930 B1	15-11-2000	
TM	438749 B	07-06-2001	
US	5922767 A	13-07-1999	
DE 19947457	A	05-04-2001	
DE	19947457 A1	05-04-2001	
AU	766365 B2	16-10-2003	
AU	7777800 A	10-05-2001	

From PCT/ISA(2) to patent family member (January 2004)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/EP 03/11134

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 19947457	A		
BR	0014465 A	11-06-2002	
CA	2385859 A1	12-04-2001	
CN	1373750 T	09-10-2002	
CZ	20021144 A3	12-06-2002	
EE	200200160 A	15-04-2003	
EP	0125189 A1	12-04-2001	
EP	1222163 A1	17-07-2002	
HR	20020264 A2	29-02-2004	
HU	0203505 A2	28-03-2003	
JP	2003511363 T	25-03-2003	
NO	20021398 A	31-05-2002	
PL	354322 A1	12-01-2004	
SK	4382002 A3	06-08-2002	
TR	200200883 T2	23-09-2002	
US	2003171351 A1	11-09-2003	
US	2004102513 A1	27-05-2004	
US	6531495 B1	11-03-2003	
ZA	200202521 A	30-10-2002	
WO 0212210	A	14-02-2002	
FR	2812876 A1	15-02-2002	
AU	8598101 A	18-02-2002	
BR	0113251 A	24-06-2003	
CA	2418895 A1	14-02-2002	
CN	1468226 T	14-01-2004	
EP	1309575 A1	14-05-2003	
WO	0212210 A1	14-02-2002	
JP	2004505958 T	26-02-2004	
US	2004039038 A1	26-02-2004	
WO 02062763	A	15-08-2002	
US	2002165394 A1	07-11-2002	
WO	02062763 A2	15-08-2002	
US 4057636	A	08-11-1977	
GB	1498679 A	26-10-1977	
AT	346357 B	10-11-1978	
AT	936075 A	15-03-1978	
AU	8712775 A	09-06-1977	
BE	836885 A1	21-06-1976	
CA	1067083 A1	27-11-1979	
CH	618429 A5	31-07-1980	
DD	122529 A5	12-10-1976	
DE	2557438 A1	24-06-1976	
DE	2560633 C2	08-09-1988	
DK	572975 A, B,	21-06-1976	
ES	443725 A1	16-04-1977	
FI	753520 A, B,	21-06-1976	
FR	2294703 A1	16-07-1976	
IT	42141 B1	04-06-1980	
IT	1062213 B	20-09-1983	
JP	1305959 C	13-03-1986	
JP	51086474 A	29-07-1976	
JP	60028819 B	06-07-1985	
LU	74071 A1	11-11-1976	
NL	7514852 A, B,	22-06-1976	
PH	12506 A	18-04-1979	
SE	424637 B	02-08-1982	
SE	7514153 A	21-06-1976	
SU	683615 A3	30-08-1979	
US	RE31244 E	17-05-1983	

From PCT/ISA(2) to patent family member (January 2004)

INTERNATIONAL SEARCH REPORT

Information on patent family members

Patent document cited in search report		Publication date		Patent family member(s)		Publication date	
US 4057636		A		YU 316275 A1 ZA 7507481 A		28-02-1982 26-01-1977	
US 5441984		A		15-08-1995		NONE	
WO 0061561		A1		19-10-2000		AU 4080300 A US 6255323 B1	
WO 0061559		A1		19-10-2000		AU 4076500 A	
WO 9518126		A1		06-07-1995		AU 1283195 A JP 9507222 T	
WO 0202534		A1		10-01-2002		AU 1675802 A	
WO 0157008		A1		09-08-2001		AU 3669801 A BG 107062 A BR 0108085 A CA 2398754 A1 CN 1422262 T EP 1254123 A1 HU 0300359 A2 JP 2003521543 T NO 20023713 A SK 12712002 A3 US 2003153568 A1	
						14-08-2001 30-04-2003 18-03-2003 09-08-2001 04-06-2003 06-11-2002 28-06-2003 15-07-2003 04-10-2002 04-02-2003 14-08-2003	

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

CORRECTED VERSION

(19) World Intellectual Property Organization
International Bureau



(10) International Publication Number
WO 2004/037789 A3

(43) International Publication Date
6 May 2004 (06.05.2004)

PCT

(51) International Patent Classification: C07D 213/68, (74) Common Representative: MERCK PATENT GMBH; 21375, 401/12, 413/12, 409/12, 417/12, A61K 31/44, 31/4427, A61P 29/00

(21) International Application Number: PCT/EP2003/011134

(22) International Filing Date: 8 October 2003 (08.10.2003)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Date: 24 October 2002 (24.10.2002) EP 02023906.7
28 July 2003 (28.07.2003) US 60490285

(71) Applicant (for all designated States except US): MERCK PATENT GMBH (DE/DE); Frankfurter Strasse 250, 64293 Darmstadt (DE).

(72) Inventors: and

(75) Inventors/Applicants (for US only): BUCHSTÄLLER, Hans-Peter (AT/DE); Heimbirgstrasse 54, 64331 Weiterstadt (DE); WIESNER, Matthias (DE/DE); Beethovenring 10, 64342 Seeheim-Jugenheim (DE); SCHADT, Oliver (DE/DE); Eschenstrasse 32, 63517 Rodenbach (DE); AMENDT, Christiane (DE/DE); Barkhausstrasse 22, 64289 Darmstadt (DE); ZENKE, Frank (DE/DE); Schulzengasse 7, 64291 Darmstadt (DE); SIRRENBERG, Christian (DE/DE); Tannstrasse 10, 64289 Darmstadt (DE); GRELL, Mathias (DE/DE); Lindenweg 44, 64291 Darmstadt (DE); FINSINGER, Dirk (DE/DE); Im Fiedlersee 5, 64291 Darmstadt (DE).

Published: with international search report

(88) Date of publication of the international search report: 28 October 2004

(48) Date of publication of this corrected version: 16 December 2004

(15) Information about Correction: see PCT Gazette No. 51/2004 of 16 December 2004, Section II

For two-letter codes and other abbreviations, refer to the "Guide to the Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHYLENE UREA DERIVATIVES AS RAF-KINASE INHIBITORS

(57) Abstract: The present invention relates to methylene urea derivatives of formula (I), the use of the compounds of formula (I) as inhibitors of raf-kinase, the use of the compounds of formula (I) for the manufacture of a pharmaceutical composition and a method of treatment, comprising administering said pharmaceutical composition to a patient.

WO 2004/037789 A3

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☒ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

THIS PAGE BLANK (USPTO)